arXiv:math/0304224v1 [math.AP] 16 Apr 2003

A DISCONTINUOUS FINITE ELEMENT APPROXIMATION OF QUASI-STATIC GROWTH OF BRITTLE FRACTURES

ALESSANDRO GIACOMINI AND MARCELLO PONSIGLIONE

Abstract. We propose a discontinuous finite element approximation for a model of quasi-static growth of brittle fractures in linearly elastic bodies formulated by Francfort and Marigo, and based on the classical Griffith's criterion. We restrict our analysis to the case of anti-planar shear and we consider discontinuous displacements which are piecewise affine with respect to a regular triangulation.

Keywords : variational models, energy minimization, free discontinuity problems, crack propagation, quasi-static evolution, brittle fracture, finite elements.

2000 Mathematics Subject Classification: 35R35, 35J25, 74R10, 35A35, 65L60.

Contents

1. Introduction 1
2. Preliminaries 5
3. The discontinuous finite element approximation 11
4. The convergence result 14
5. Piecewise Affine Transfer of Jump and Proof of Proposition 4.2 23
6. Revisiting the approximation by Francfort and Larsen 32
References 34

1. Introduction

In this paper we formulate a discontinuous finite element approximation for a model of quasi-static growth of brittle fractures in linearly elastic bodies proposed by Francfort and Marigo [11]. Their model is based on the classical Griffith's criterion which involves a competition between *bulk* and *surface* energies. To be precise, let $\Omega \subseteq \mathbb{R}^3$ be an elastic body, $\partial_D\Omega$ be a part of its boundary and let $g : \partial_D\Omega \to \mathbb{R}^3$ be the spatial displacement of Ω at the points of $\partial_D\Omega$. According to Griffith's theory, given a preexisting crack $\Gamma_1 \subseteq \overline{\Omega}$, the new crack Γ and the displacement $u : \Omega \setminus \Gamma \to \mathbb{R}^3$ associated to g at the equilibrium minimizes the following energy

$$\mathcal{E}(v, g, \Gamma) := \int_\Omega \mu |Ev|^2 + \frac{\lambda}{2} |\mathrm{div}\, v|^2 \, dx + \mathcal{H}^2(\Gamma), \tag{1.1}$$

among all cracks Γ with $\Gamma_1 \subseteq \Gamma$ and all displacements $v : \Omega \setminus \Gamma \to \mathbb{R}^3$ with $v = g$ on $\partial_D\Omega \setminus \Gamma$. Here Ev denotes the symmetric part of the gradient of v, and $\mathcal{H}^2$ denotes the two dimensional Hausdorff measure, while μ and λ are the Lamé coefficients. The boundary condition is required only on $\partial_D\Omega \setminus \Gamma$ because the displacement in a fractured region is assumed not to be transmitted. Let us indicate by $\mathcal{E}(g,\Gamma)$ the minimum value of (1.1) among all $v : \Omega \setminus \Gamma \to \mathbb{R}^3$ with $v = g$ on $\partial_D\Omega \setminus \Gamma$.

Supposing that the boundary displacement g varies with the time $t \in [0,1]$, the quasi-static evolution $t \to \Gamma(t)$ proposed in [11] requires that:

(1) $\Gamma(t)$ is increasing in time, i.e., $\Gamma(t_1) \subseteq \Gamma(t_2)$ for all $0 \le t_1 \le t_2 \le 1$ (irreversibility of the process);

(2) $\mathcal{E}(g(t),\Gamma(t)) \le \mathcal{E}(g(t),\Gamma)$ for all cracks Γ such that $\cup_{s<t}\Gamma(s) \subseteq \Gamma$ (equilibrium condition);

(3) the total energy $\mathcal{E}(g(t),\Gamma(t))$ is absolutely continuous in time and (conservation of energy)

$$\frac{d}{dt}\mathcal{E}(g(t),\Gamma(t)) = 2\mu \int_\Omega Eu(t)E\dot{g}(t)\,dx + \lambda \int_\Omega \operatorname{div} u(t) \operatorname{div} \dot{g}(t)\,dx.$$

A precise mathematical formulation of this model has been given by Dal Maso and Toader [9] in the case of *anti-planar shear* in dimension two assuming that the fractures are compact sets with a finite number of connected components. Recently Francfort and Larsen [10], using the framework of SBV-functions, proved the existence of a quasi-static growth of brittle fractures in the case of *generalized anti-planar shear* and without assumptions on the structure of the fractures which are dealt with the set of jumps of the displacements. To be precise, they consider as elastic body an infinite cylinder whose section $\Omega \subseteq \mathbb{R}^N$ is subject to a displacement $u \in SBV(\Omega)$ in the direction orthogonal to Ω. The crack at time t on the section Ω is defined as

$$\Gamma(t) := \bigcup_{s<t} \left[S_{u(s)} \cup (\partial_D\Omega \cap \{u(s) \neq g(s)\}) \right],$$

where S_u denotes the set of jumps of u. Moreover the pair $(u(t),\Gamma(t))$ is such that:

(a) for all $v \in SBV(\Omega)$

$$\int_\Omega |\nabla u(t)|^2\,dx + \mathcal{H}^{N-1}(\Gamma(t)) \le \int_\Omega |\nabla v|^2\,dx + \mathcal{H}^{N-1}(S_v \cup (\partial_D\Omega \cap \{v \neq g(t)\}) \cup \Gamma(t)); \tag{1.2}$$

(b) the total energy $\mathcal{E}(t) := \int_\Omega |\nabla u(t)|^2\,dx + \mathcal{H}^{N-1}(\Gamma(t))$ is absolutely continuous and

$$\mathcal{E}(t) = \mathcal{E}(0) + 2\int_0^t \int_\Omega \nabla u(\tau)\nabla\dot{g}(\tau)\,dx\,d\tau. \tag{1.3}$$

The aim of this paper is to discretize the model using a suitable finite element method and to a give a rigorous proof of its convergence to a quasi-static evolution in the sense of Francfort and Larsen. We restrict our analysis to a two dimensional setting considering only a polygonal reference configuration $\Omega \subseteq \mathbb{R}^2$.

The discretization of the domain Ω is carried out, following [12], considering two parameters $\varepsilon > 0$ and $a \in]0, \frac{1}{2}[$. We consider a regular triangulation $\mathbf{R}_\varepsilon$ of size ε of Ω, i.e. we assume that there exist two constants c_1 and c_2 so that every triangle $T \in \mathbf{R}_\varepsilon$ contains a ball of radius $c_1\varepsilon$ and is contained in a ball of radius $c_2\varepsilon$. In order to treat the boundary data, we assume also that $\partial_D\Omega$ is composed of edges of $\mathbf{R}_\varepsilon$. On each edge $[x,y]$ of $\mathbf{R}_\varepsilon$ we consider a point z such that $z = tx + (1-t)y$ with $t \in [a, 1-a]$. These points are called *adaptive vertices*. Connecting together the adaptive vertices, we divide every $T \in \mathbf{R}_\varepsilon$ into four triangles. We take the new triangulation $\mathbf{T}$ obtained after this division as the discretization of Ω. The family of all such triangulations is denoted by $\mathcal{T}_{\varepsilon,a}(\Omega)$.

The discretization of the energy functional is obtained restricting the total energy to the family of functions u which are affine on the triangles of some triangulation $\mathbf{T}(u) \in \mathcal{T}_{\varepsilon,a}(\Omega)$ and are allowed to jump across the edges of $\mathbf{T}(u)$. We indicate this space by $\mathcal{A}_{\varepsilon,a}(\Omega)$. The boundary data is assumed to belong to the space $\mathcal{AF}_\varepsilon(\Omega)$ of continuous functions which are affine on every triangle $T \in \mathbf{R}_\varepsilon$.

Given the boundary data $g \in W^{1,1}([0,1], H^1(\Omega))$ with $g(t) \in \mathcal{AF}_\varepsilon(\Omega)$ for all $t \in [0,1]$, we divide $[0,1]$ into subintervals $[t_i^\delta, t_{i+1}^\delta]$ of size $\delta > 0$ for $i = 0, \dots, N_\delta$, we set $g_i^\delta = g(t_i^\delta)$, and for all $u \in \mathcal{A}_{\varepsilon,a}(\Omega)$ we indicate by $S_D^{g_i^\delta}(u)$ the edges of the triangulation $\mathbf{T}(u)$ contained in $\partial_D\Omega$ on which $u \neq g_i^\delta$. Using a variational argument we construct a *discrete evolution* $\{u_{\varepsilon,a}^{\delta,i} : i = 0, \dots, N_\delta\}$ such that $u_{\varepsilon,a}^{\delta,i} \in \mathcal{A}_{\varepsilon,a}(\Omega)$ for all $i = 0, \dots, N_\delta$, and such that considering the *discrete fracture*

$$\Gamma_{\varepsilon,a}^{\delta,i} := \bigcup_{r=0}^{i} \left[S_{u_{\varepsilon,a}^{\delta,r}} \cup S_D^{g_r^\delta}(u_{\varepsilon,a}^{\delta,r}) \right],$$

the following *unilateral minimality property* holds:

$$\int_\Omega |\nabla u_{\varepsilon,a}^{\delta,i}|^2 \, dx \leq \int_\Omega |\nabla v|^2 \, dx + \mathcal{H}^1 \left(\left(S_v \cup S_D^{g_i^\delta}(v) \right) \setminus \Gamma_{\varepsilon,a}^{\delta,i-1} \right). \tag{1.4}$$

Moreover we get suitable estimates for the discrete total energy

$$\mathcal{E}_{\varepsilon,a}^{\delta,i} := \|\nabla u_{\varepsilon,a}^{\delta,i}\|^2_{L^2(\Omega;\mathbb{R}^2)} + \mathcal{H}^1 \left(\Gamma_{\varepsilon,a}^{\delta,i} \right).$$

The definition of the discrete fracture ensures that $\Gamma_{\varepsilon,a}^{\delta,i} \subseteq \Gamma_{\varepsilon,a}^{\delta,j}$ for all $i \leq j$, recovering in this discrete setting the irreversibilty of the growth given in (1). The minimality property (1.4) is the reformulation in the finite element space of the equilibrium condition (2).

In order to perform the asymptotic analysis of the *discrete evolution* $\{u_{\varepsilon,a}^{\delta,i} : i = 0, \dots, N_\delta\}$, we make the piecewise constant interpolation in time $u_{\varepsilon,a}^\delta(t) = u_{\varepsilon,a}^{\delta,i}$ and $\Gamma_{\varepsilon,a}^\delta(t) = \Gamma_{\varepsilon,a}^{\delta,i}$ for all $t_i^\delta \leq t < t_{i+1}^\delta$. The main result of the paper is the following theorem.

Theorem 1.1. *Let* $g \in W^{1,1}([0,1], H^1(\Omega))$ *be such that* $\|g(t)\|_\infty \leq C$ *for all* $t \in [0,1]$ *and let* $g_\varepsilon \in W^{1,1}([0,1], H^1(\Omega))$ *be such that* $\|g_\varepsilon(t)\|_\infty \leq C$, $g_\varepsilon(t) \in \mathcal{AF}_\varepsilon(\Omega)$ *for all* $t \in [0,1]$ *and*

$$g_\varepsilon \to g \quad \text{strongly in } W^{1,1}([0,1], H^1(\Omega)). \tag{1.5}$$

Given the discrete evolution $\{t \to u_{\varepsilon,a}^\delta(t)\}$ *relative to the boundary data* g_ε, *let* $\Gamma_{\varepsilon,a}^\delta$ *and* $\mathcal{E}_{\varepsilon,a}^\delta$ *be the associated fracture and total energy.*

Then there exist $\delta_n \to 0$, $\varepsilon_n \to 0$, $a_n \to 0$, *and a quasi-static evolution* $\{t \to (u(t), \Gamma(t)),\ t \in [0,1]\}$ *relative to the boundary data* g, *satisfying* (1.2) *and* (1.3), *and such that setting* $u_n := u_{\varepsilon_n,a_n}^{\delta_n}$, $\Gamma_n := \Gamma_{\varepsilon_n,a_n}^{\delta_n}$, $\mathcal{E}_n := \mathcal{E}_{\varepsilon_n,a_n}^{\delta_n}$, *the following hold:*

(a) *if* $\mathcal{N}$ *is the set of discontinuities of* $\mathcal{H}^1(\Gamma(\cdot))$, *for all* $t \in [0,1] \setminus \mathcal{N}$ *we have*

$$\nabla u_n(t) \to \nabla u(t) \quad \text{strongly in } L^2(\Omega; \mathbb{R}^2) \tag{1.6}$$

and

$$\lim_n \mathcal{H}^1(\Gamma_n(t)) = \mathcal{H}^1(\Gamma(t)); \tag{1.7}$$

(b) *for all* $t \in [0,1]$ *we have*

$$\lim_n \mathcal{E}_n(t) = \mathcal{E}(t). \tag{1.8}$$

We conclude that we have the convergence of the total energy at each time $t \in [0,1]$, and the separate convergence of bulk and surface energy for all $t \in [0,1]$ except a countable set.

In order to prove Theorem 1.1, we proceed in two steps. Firstly, we fix a and let $\delta \to 0$ and $\varepsilon \to 0$. We obtain an evolution $\{t \to u_a(t) \,:\, t \in [0,1]\}$ such that $\nabla u^\delta_{\varepsilon,a}(t) \to \nabla u_a(t)$ strongly in $L^2(\Omega;\mathbb{R}^2)$ for all t up to a countable set and such that the following minimality property holds: for all $v \in SBV(\Omega)$

$$\int_\Omega |\nabla u_a(t)|^2\,dx \le \int_\Omega |\nabla v|^2\,dx + \mu(a)\mathcal{H}^1\left(\left(S_v \cup (\partial_D\Omega \cap \{v \neq g(t)\})\right) \setminus \Gamma_a(t)\right), \tag{1.9}$$

where $\mu :]0,\frac{1}{2}[\to]0,+\infty[$ is a function independent of ε and δ, such that $\mu \ge 1$, $\lim_{a\to 0}\mu(a) = 1$ and $\Gamma_a(t) := \bigcup_{s \le t, s \in D} S_{u_a(s)} \cup (\partial_D\Omega \cap \{u_a(s) \neq g(s)\})$. The minimality property (1.9) takes into account possible anisotropies that could be generated as δ and $\varepsilon \to 0$: in fact, since a is fixed, we have that the angles of the triangles in $\mathcal{T}_{\varepsilon,a}(\Omega)$ are between fixed values (determined by a), and so fractures with certain directions cannot be approximated in length. In the second step, we let $a \to 0$ and determine from $\{t \to u_a(t) \,:\, t \in [0,1]\}$ a quasi-static evolution $\{t \to u(t) \,:\, t \in [0,1]\}$ in the sense of Francfort and Larsen. Then, using a diagonal argument, we find sequences $\delta_n \to 0$, $\varepsilon_n \to 0$, and $a_n \to 0$ satisfying Theorem 1.1.

The main difficulties arise in the first part of our analysis, namely when $\delta, \varepsilon \to 0$. The convergence $u^\delta_{\varepsilon,a}(t) \to u_a(t)$ in $SBV(\Omega)$ for $t \in D \subseteq [0,1]$ countable and dense is easily obtained by means of Ambrosio's Compactness Theorem. The minimality property (1.9) derives from its discrete version (1.4) using a variant of Lemma 1.2 of [10]: given $v \in SBV(\Omega)$, we construct $v^\delta_{\varepsilon,a} \in \mathcal{A}_{\varepsilon,a}(\Omega)$ such that

$$\nabla v^\delta_{\varepsilon,a} \to \nabla v \quad \text{strongly in } L^2(\Omega;\mathbb{R}^2) \tag{1.10}$$

and

$$\limsup_{\delta,\varepsilon\to 0} \mathcal{H}^1\left[\left(S_{v^\delta_{\varepsilon,a}} \cup S_D^{g^\delta_\varepsilon(t)}(v^\delta_{\varepsilon,a})\right) \setminus \Gamma^\delta_{\varepsilon,a}(t)\right] \le \tag{1.11}$$
$$\le \mu(a)\mathcal{H}^1\left[\left(S_v \cup \left(\partial_D\Omega \cap \{v \neq g(t)\}\right)\right) \setminus \Gamma_a(t)\right],$$

where $g^\delta_\varepsilon(t) := g_\varepsilon(t^\delta_i)$ for $t^\delta_i \le t < t^\delta_{i+1}$. The main difference with respect to Lemma 1.2 of [10] is that we have to find the approximating functions $v^\delta_{\varepsilon,a}$ in the finite element space $\mathcal{A}_{\varepsilon,a}(\Omega)$. This can be regarded as an interpolation problem, so we try to construct triangulations $\mathbf{T}_\varepsilon \in \mathcal{T}_{\varepsilon,a}(\Omega)$ adapted to v in order to obtain (1.10) and (1.11). In all the geometric operations involved, we need to avoid degeneration of the triangles of $\mathbf{T}(u^\delta_{\varepsilon,a}(t))$ which is guaranteed from the fact that a is constant: this is the principal reason to keep a fixed in the first step. A second difficulty arises when $u_a(\cdot)$ is extended from D to the entire interval $[0,1]$: indeed it is no longer clear whether $\nabla u^\delta_{\varepsilon,a}(t) \to \nabla u_a(t)$ for $t \notin D$. Since the space $\mathcal{A}_{\varepsilon,a}(\Omega)$ is not a vector space, we cannot provide an estimate on $\|\nabla u^\delta_{\varepsilon,a}(t) - \nabla u^\delta_{\varepsilon,a}(s)\|$ with $s \in D$ and $s < t$: we thus cannot expect to recover the convergence at time t from the convergence at time s. We overcome this difficulty observing that $\nabla u^\delta_{\varepsilon,a}(t) \to \nabla \tilde{u}_a$ with $\tilde{u}_a$ satisfying a minimality property similar to (1.9) and then proving $\nabla \tilde{u}_a = \nabla u_a(t)$ by a uniqueness argument for the gradients of the solutions.

The plan of the paper is the following. In Section 2 we give the basic definitions and prove some auxiliary results. In Section 3, we prove the existence of a discrete evolution. In Section 4 we prove the convergence of the discrete evolution to a quasi-static evolution of brittle fractures in the sense of Francfort and Larsen. The proof of minimality property (1.9) requires a careful analysis to which is dedicated Section 5. In Section 6 we show that the arguments of Section 4 can be used to improve the convergence results for the discrete in time approximation considered in [10].

2. Preliminaries

In this section we state the notation and prove some preliminaries employed in the rest of the paper.

Basic notation. We will employ the following basic notation:

- Ω is a polygonal open subset of $\mathbb{R}^2$;
- $L^p(\Omega;\mathbb{R}^m)$ with $1 \le p < +\infty$ and $m \ge 1$ is the usual Lebesgue space of p-summable $\mathbb{R}^m$-valued functions, and $L^p(\Omega) := L^p(\Omega;\mathbb{R})$;
- for all $k \ge 1$ and $1 \le p \le +\infty$, $W^{k,p}(\Omega)$ is the usual Sobolev space of functions in $L^p(\Omega)$ with distributional derivatives of order $1,\dots,k$ in $L^p(\Omega)$; we will write $H^k(\Omega)$ for $W^{k,2}(\Omega)$;
- if $u \in W^{k,p}(\Omega)$, ∇u is its gradient;
- $\mathcal{H}^1$ is the one-dimensional Hausdorff measure;
- $\|\cdot\|_\infty$ denotes the sup-norm;
- if $f \in L^2(\Omega;\mathbb{R}^m)$, $\|f\|$ denotes the L^2-norm of f;
- for all $A \subseteq \mathbb{R}^2$, $|A|$ denotes the Lebesgue measure of A;
- if μ is a measure on $\mathbb{R}^2$ and A is a Borel subset of $\mathbb{R}^2$, $\mu\llcorner A$ denotes the restriction of μ to A, i.e. $(\mu\llcorner A)(B) := \mu(B\cap A)$ for all Borel sets $B \subseteq \mathbb{R}^2$;
- if $\sigma \in]0,+\infty[$, o_σ is such that $\lim_{\sigma\to 0^+} o_\sigma = 0$.

Special functions of bounded variation. For the general theory of functions of bounded variation, we refer to [4]; here we recall some basic definitions and theorems we need in the sequel. Let A be an open subset of $\mathbb{R}^N$, and let $u : A \to \mathbb{R}^n$. We say that $u \in BV(A;\mathbb{R}^n)$ if $u \in L^1(A;\mathbb{R}^n)$, and its distributional derivative is a vector-valued Radon measure on A. We say that $u \in SBV(A;\mathbb{R}^n)$ if $u \in BV(A;\mathbb{R}^n)$ and its distributional derivative can be represented as

$$Du(A) = \int_A \nabla u(x)\,dx + \int_{A\cap S_u} (u^+(x) - u^-(x)) \otimes \nu_x \, d\mathcal{H}^{N-1}(x),$$

where ∇u denotes the approximate gradient of u, S_u denotes the set of approximate jumps of u, u^+ and u^- are the traces of u on S_u, ν_x is the normal to S_u at x, and $\mathcal{H}^{N-1}$ is the $(N-1)$-dimensional Hausdorff measure. The space $SBV(A;\mathbb{R}^n)$ is called the space of *special functions of bounded variation.* Note that if $u \in SBV(A;\mathbb{R}^n)$, then the singular part of Du is concentrated on S_u which turns out to be countably $\mathcal{H}^{N-1}$-rectifiable.

The space SBV is very useful when dealing with variational problems involving volume and surface energies because of the following compactness and lower semicontinuity result due to L.Ambrosio (see [1], [2], [3], [4]).

Theorem 2.1. *Let A be an open and bounded subset of $\mathbb{R}^N$, and let (u_k) be a sequence in $SBV(A;\mathbb{R}^n)$. Assume that there exists $q>1$ and $c\geq 0$ such that*

$$\int_A |\nabla u_k|^q\,dx+\mathcal{H}^{N-1}(S_{u_k})+||u_k||_\infty\leq c$$

for every $k\in\mathbb{N}$. Then there exists a subsequence (u_{k_h}) and a function $u\in SBV(A;\mathbb{R}^n)$ such that

$$\begin{aligned} &u_{k_h}\to u \quad \textit{strongly in } L^1(A;\mathbb{R}^n);\\ &\nabla u_{k_h}\rightharpoonup \nabla u \quad \textit{weakly in } L^1(A;M^{N\times n});\\ &\mathcal{H}^{N-1}(S_u)\leq \liminf_h \mathcal{H}^{N-1}(S_{u_{k_h}}). \end{aligned} \tag{2.1}$$

In the rest of the paper, we will say that $u_k\to u$ in $SBV(A;\mathbb{R}^n)$ if u_k and u satisfy (2.1). It will also be useful the following fact which can be derived from Ambrosio's Theorem: if $u_k\to u$ in $SBV(A;\mathbb{R}^n)$ and if $\mathcal{H}^{N-1}\llcorner S_{u_k}\overset{*}{\rightharpoonup}\mu$ weakly-star in the sense of measures, then $\mathcal{H}^{N-1}\llcorner S_u\leq\mu$ as measures. We will set $SBV(A):=SBV(A;\mathbb{R})$.

Quasi-static evolution of brittle fracture. Let Ω be an open bounded subset of $\mathbb{R}^N$ with Lipschitz boundary, and let $\partial_D\Omega$ be a subset of $\partial\Omega$ open in the relative topology. Let $g:[0,1]\to H^1(\Omega)$ be absolutely continuous; we indicate the gradient of g at time t by $\nabla g(t)$, and the time derivative of g at time t by $\dot g(t)$. The main result of [10] is the following theorem.

Theorem 2.2. *There exists a crack $\Gamma(t)\subseteq\overline{\Omega}$ and a field $u(t)\in SBV(\Omega)$ such that:*

(a) *$\Gamma(t)$ increases with t;*

(b) *$u(0)$ minimizes*

$$\int_\Omega |\nabla v|^2\,dx+\mathcal{H}^{N-1}(S_v\cup\{x\in\partial_D\Omega : v(x)\neq g(0)(x)\})$$

among all $v\in SBV(\Omega)$ (inequalities on $\partial_D\Omega$ are intended for the traces of v and g);

(c) *for $t>0$, $u(t)$ minimizes*

$$\int_\Omega |\nabla v|^2\,dx+\mathcal{H}^{N-1}\left([S_v\cup\{x\in\partial_D\Omega : v(x)\neq g(t)(x)\}]\setminus\Gamma(t)\right)$$

among all $v\in SBV(\Omega)$;

(d) *$S_{u(t)}\cup\{x\in\partial_D\Omega : u(t)(x)\neq g(t)(x)\}\subseteq\Gamma(t)$, up to a set of $\mathcal{H}^{N-1}$-measure 0.*

Furthermore, the total energy

$$\mathcal{E}(t):=\int_\Omega |\nabla u(t)|^2\,dx+\mathcal{H}^{N-1}(\Gamma(t))$$

is absolutely continuous and satisfies

$$\mathcal{E}(t)=\mathcal{E}(0)+2\int_0^t\int_\Omega \nabla u(\tau)\nabla\dot g(\tau)\,dx\,d\tau$$

for every $t \in [0,1]$. *Finally, for any countable, dense set* $I \subseteq [0,1]$, *the crack* $\Gamma(t)$ *and the displacement* $u(t)$ *can be chosen so that*

$$\Gamma(t) = \bigcup_{\tau \in I, \tau \leq t} \left(S_{u(\tau)} \cup \{x \in \partial_D \Omega \,:\, u(\tau)(x) \neq g(\tau)(x)\} \right).$$

The main tool in the proof of Theorem 2.2 is the following result [10, Theorem 2.1], which is useful also in our analysis.

Theorem 2.3. *Let* $\overline{\Omega} \subseteq \Omega'$, *with* $\partial\Omega$ *Lipschitz, and let for* $r = 1, \dots, i$ (u_n^r) *be a sequence in* $SBV(\Omega')$ *such that*

(1) $S_{u_n^r} \subseteq \overline{\Omega}$;

(2) $|\nabla u_n^r|$ *weakly converges in* $L^1(\Omega')$; *and*

(3) $u_n^r \to u^r$ *strongly in* $L^1(\Omega')$,

where $u^r \in BV(\Omega')$ *with* $\mathcal{H}^{N-1}(S_{u^r}) < \infty$. *Then for every* $\phi \in SBV(\Omega')$ *with* $\mathcal{H}^{N-1}(S_\phi) < \infty$ *and* $\nabla\phi \in L^q(\Omega; \mathbb{R}^N)$ *for some* $q \in [1, +\infty[$, *there exists a sequence* (ϕ_n) *in* $SBV(\Omega')$ *with* $\phi_n \equiv \phi$ *on* $\Omega' \setminus \overline{\Omega}$ *such that*

(a) $\phi_n \to \phi$ *strongly in* $L^1(\Omega')$;

(b) $\nabla\phi_n \to \nabla\phi$ *strongly in* $L^q(\Omega')$; *and*

(c) $\mathcal{H}^{N-1}\left([S_{\phi_n} \setminus \bigcup_{r=1}^i S_{u_n^r}] \setminus [S_\phi \setminus \bigcup_{r=1}^i S_{u^r}] \right) \to 0.$

In particular

$$\limsup_n \mathcal{H}^{N-1}\left(S_{\phi_n} \setminus \bigcup_{r=1}^i S_{u_n^r} \right) \leq \mathcal{H}^{N-1}\left(S_\phi \setminus \bigcup_{r=1}^i S_{u^r} \right). \tag{2.2}$$

Hausdorff metric on compact sets. Let $A \subseteq \mathbb{R}^2$ be open and bounded, and let $\mathcal{K}(\overline{A})$ be the set of all compact subsets of $\overline{A}$. $\mathcal{K}(\overline{A})$ can be endowed by the Hausdorff metric d_H defined by

$$d_H(K_1, K_2) := \max\left\{ \sup_{x \in K_1} \operatorname{dist}(x, K_2), \sup_{y \in K_2} \operatorname{dist}(y, K_1) \right\},$$

with the conventions $\operatorname{dist}(x, \emptyset) = \operatorname{diam}(A)$ and $\sup \emptyset = 0$, so that $d_H(\emptyset, K) = 0$ if $K = \emptyset$ and $d_H(\emptyset, K) = \operatorname{diam}(A)$ if $K \neq \emptyset$. It turns out that $\mathcal{K}(\overline{A})$ endowed with the Hausdorff metric is a compact space (see e.g. [13]).

Triangulations. Let $\Omega \subseteq \mathbb{R}^2$ be a polygonal set and let us fix two positive constants $0 < c_1 < c_2$. By a *regular triangulation* of Ω of size ε we intend a finite family of (closed) triangles T_i such that $\overline{\Omega} = \bigcup_i T_i$, $T_i \cap T_j$ is either empty or equal to a common edge or to a common vertex, and each T_i contains a ball of diameter $c_1\varepsilon$ and is contained in a ball of diameter $c_2\varepsilon$.

We indicate by $\mathcal{R}_\varepsilon(\Omega)$ the family of all regular triangulations of Ω of size ε. It turns out that there exist $0 < \vartheta_1 < \vartheta_2 < \pi$ such that for all T belonging to a triangulation $\mathbf{T} \in \mathcal{R}_\varepsilon(\Omega)$, the inner angles of T are between ϑ_1 and ϑ_2. Moreover, every edge of T has length greater than $c_1\varepsilon$ and lower than $c_2\varepsilon$.

Let us fix a triangulation $\mathbf{R}_\varepsilon \in \mathcal{R}_\varepsilon(\Omega)$ for all $\varepsilon > 0$ and let $a \in]0, \frac{1}{2}[$. Let us consider a new triangulation $\mathbf{T}$ nested in $\mathbf{R}_\varepsilon$ obtained dividing each $T \in \mathbf{R}_\varepsilon$ into four triangles taking over every edge $[x, y]$ of T a knot z which satisfies

$$z = tx + (1-t)y, \qquad t \in [a, 1-a].$$

We will call these new vertices *adaptive*, the triangles obtained joining these points *adaptive triangles*, and their edges *adaptive edges* (see Fig.1).



Fig. 1

We denote by $\mathcal{T}_{\varepsilon,a}(\Omega)$ the set of all triangulations $\mathbf{T}$ constructed in this way. Note that for all $\mathbf{T} \in \mathcal{T}_{\varepsilon,a}(\Omega)$ there exists $0 < c_1^a < c_2^a < +\infty$ such that every $T_i \in \mathbf{T}$ contains a ball of diameter $c_1^a\varepsilon$ and is contained in a ball of diameter $c_2^a\varepsilon$. Then there exist $0 < \vartheta_1^a < \vartheta_2^a < \pi$ such that for all triangles T belonging to a triangulation $\mathbf{T} \in \mathcal{T}_{\varepsilon,a}(\Omega)$, the inner angles of T are between ϑ_1^a and ϑ_2^a. Moreover, every edge of T has length greater than $c_1^a\varepsilon$ and lower than $c_2^a\varepsilon$.

We will often use the following *interpolation estimate* (see [7, Theorem 3.1.5]). If $u \in W^{2,2}(\Omega)$ and $T \in \mathbf{R}_\varepsilon$, let u_T denote the affine interpolation of u on T. We have that there exists K depending only on c_1, c_2 such that

$$\|u_T - u\|_{W^{1,2}(T)} \le K\varepsilon \|u\|_{W^{2,2}(T)}. \tag{2.3}$$

Estimate (2.3) holds also for $\mathbf{T} \in \mathcal{T}_{\varepsilon,a}(\Omega)$: in this case K depends on a.

Some elementary constructions. The following lemmas will be used in Section 4.

Lemma 2.4. *Let $\mathbf{T} \in \mathcal{T}_{\varepsilon,a}(\Omega)$, and let $l \subseteq \Omega$ be a segment with extremes p, q belonging to edges of $\mathbf{T}$. There exists a polyhedral curve Γ with extremal points p and q (see Fig.2) such* *that* Γ *is contained in the union of the edges of those* $T \in \mathbf{T}$ *with* $T \cap l \neq \emptyset$*, and such that the following properties hold:*

(1) $\Gamma = \gamma_p \cup \gamma \cup \gamma_q$*, where* γ *is union of edges of* $\mathbf{T}$ *and* γ_p*,* γ_q *are segments containing* p *and* q *respectively, and each one is contained in an edge of* $\mathbf{T}$*;*

(2) *there exists a constant* c *independent of* ϵ *(but depending on* a*) such that*

$$\mathcal{H}^1(\Gamma) \leq c\,\mathcal{H}^1(l).$$



Fig. 2

Proof. Let $\{T_1, \dots T_k\}$ be the family of triangles in $\mathbf{T}$ such that the intersection with l is a segment with positive length. For every integer $1 \leq i \leq k$, let $l_i := T_i \cap l$. If l_i is an edge of T_i, we set $D_i = T_i$. Otherwise let D_i be a connected component of $T_i \setminus l_i$ such that $|D_i| \leq \frac{1}{2}|T_i|$. We claim that there exists a constant $c > 0$ independent of ϵ such that

$$\mathcal{H}^1(\partial D_i) \leq c\,\mathcal{H}^1(l_i). \tag{2.4}$$

We have to analyze two possibilities, namely D_i is a triangle, or D_i is a trapezoid. Suppose that D_i is a triangle and that m_i is an edge of D_i. Let α be the angle of D_i opposite to l_i. It is easy to prove that $\mathcal{H}^1(l_i) \geq \mathcal{H}^1(m_i)\sin\alpha$, and so

$$\mathcal{H}^1(l_i) \geq \frac{1}{3}\sin\alpha\mathcal{H}^1(\partial D_i).$$

Since $\vartheta_1^a \leq \alpha \leq \vartheta_2^a$, $\sin\alpha$ is uniformly bounded from below, and hence inequality (2.4) follows. If D_i is a trapezoid, since $|D_i| \leq \frac{1}{2}|T_i|$, it follows that $T_i \setminus D_i$ is a triangle such that its edges different from l_i have length greater than $\frac{1}{2}c_1^a\varepsilon$. Let α be the inner angle of $T_i \setminus D_i$ opposite to l_i. We have that

$$\mathcal{H}^1(l_i) \geq \frac{1}{2}\sin\alpha c_1^a\varepsilon \geq \frac{1}{2}\sin\alpha\frac{c_1^a}{c_2^a}\frac{1}{4}\mathcal{H}^1(\partial D_i).$$

Since $\vartheta_1^a \leq \alpha \leq \vartheta_2^a$, inequality (2.4) follows.

By (2.4), we deduce that

$$\mathcal{H}^1\left(\bigcup_{i=1}^{k}\partial D_i\right) \le c\,\mathcal{H}^1(l);$$

moreover, since $\bigcup_{i=1}^{k}\partial D_i$ is arcwise connected and contains p, q, we conclude that there exists a curve $\Gamma \subseteq \bigcup_{i=1}^{k}\partial D_i$ which satisfies the thesis. □

Lemma 2.5. *There exists a constant $c > 0$ such that for every segment $l \subseteq \Omega$ there exists ε_0 with the following property: for every $\varepsilon \le \varepsilon_0$, setting $\mathcal{R}(l) := \{T \in \mathbf{R}_\varepsilon : T \cap l \neq \emptyset\}$, we have*

$$\mathcal{H}^1(\partial\mathcal{R}(l)) \le c\mathcal{H}^1(l).$$

Proof. Let $\mathcal{N}_\varepsilon(l) := \{x \in \Omega : \operatorname{dist}(x, l) \le c_2\varepsilon\}$. We have that $|\mathcal{N}_\varepsilon(l)| = \mathcal{H}^1(l)c_2\varepsilon + \pi c_2^2\varepsilon^2$, and hence there exists a positive constant ε_0 such that, for every $\varepsilon \le \varepsilon_0$, we have that

$$|\mathcal{N}_\varepsilon(l)| \le 2\mathcal{H}^1(l)c_2\varepsilon.$$

We have that $\mathcal{R}(l) \subseteq \mathcal{N}_\varepsilon(l)$, and

$$\sharp\mathcal{R}(l) \le \frac{4}{c_1^2\pi^2}\frac{|\mathcal{N}_\varepsilon(l)|}{\varepsilon^2},$$

where $\sharp\mathcal{R}(l)$ denotes the number of triangles of $\mathcal{R}(l)$. Then, we have

$$\mathcal{H}^1(\partial\mathcal{R}(l)) \le 3c_2\varepsilon\,\sharp\mathcal{R}(l) \le 3c_2\varepsilon\frac{4}{c_1^2\pi^2}\frac{|\mathcal{N}_\varepsilon(l)|}{\varepsilon^2} \le 3c_2^2\frac{4}{c_1^2\pi^2}2\mathcal{H}^1(l),$$

and so the proof is concluded. □

A density result. Let $A \subseteq \mathbb{R}^2$ be open. We say that $K \subseteq A$ is polygonal (with respect to A), if it is the intersection of A with the union of a finite number of closed segments. The following density result is proved in [8].

Theorem 2.6. *Assume that ∂A is locally Lipschitz, and let $u \in SBV(A)$ such that $u \in L^2(A)$, $\nabla u \in L^2(\Omega;\mathbb{R}^2)$, and $\mathcal{H}^1(S_u) < +\infty$. For every $\varepsilon > 0$, there exists a function $v \in SBV(A)$ such that*

(a) *S_v is essentially closed, i.e., $\mathcal{H}^1(\overline{S_v} \setminus S_v) = 0$;*

(b) *$\overline{S_v}$ is a polyhedral set;*

(c) *$v \in W^{k,\infty}(A \setminus \overline{S_v})$ for every $k \in \mathbb{N}$;*

(d) *$||v - u||_{L^2(A)} < \varepsilon$;*

(e) *$||\nabla v - \nabla u||_{L^2(A;\mathbb{R}^2)} < \varepsilon$;*

(f) *$|\mathcal{H}^1(S_v) - \mathcal{H}^1(S_u)| < \varepsilon$.*

Let $\partial_D\Omega$ be a relatively open subset of $\partial\Omega$ composed of edges lying in $\partial\Omega$. Let us consider Ω_D polygonal open bounded subset of $\mathbb{R}^2$ such that $\Omega_D \cap \Omega = \emptyset$ and $\partial\Omega \cap \partial\Omega_D = \partial_D\Omega$ up to a finite number of vertices. We set $\Omega' := \Omega \cup \Omega_D \cup \partial_D\Omega$. In Section 4, we will use the following result.

Proposition 2.7. *Given $u \in SBV(\Omega')$ with $u = 0$ on $\Omega' \setminus \overline{\Omega}$ and $\mathcal{H}^{N-1}(S_u) < +\infty$, there exists $u_h \in SBV(\Omega')$ such that*

(a) *$u_h = 0$ in $\Omega' \setminus \overline{\Omega}$;*

(b) *S_{u_h} is polyhedral, $\overline{S_{u_h}} \subseteq \Omega$ and $u_h \in W^{k,\infty}(\Omega' \setminus \overline{S_{u_h}})$ for all k;*

(c) *$u_h \to u$ strongly in $L^2(\Omega')$ and $\nabla u_h \to \nabla u$ strongly in $L^2(\Omega'; \mathbb{R}^2)$;*

(d) *for all A open subset of Ω' with $\mathcal{H}^1(\partial A \cap S_u) = 0$, we have*

$$\lim_h \mathcal{H}^1(A \cap S_{u_h}) = \mathcal{H}^1(A \cap S_u).$$

Proof. Using a partition of unity, we may prove the result in the case $\Omega :=]-1,1[\times]0,1[$, $\Omega' :=]-1,1[\times]-1,1[$, and $\partial_D\Omega :=]-1,1[\times\{0\}$. We set $w_h(x,y) := u(x, y-h)$, and let φ_h be a cut off function with $\varphi_h = 1$ on $]-1,1[\times]-1,\frac{h}{3}[$, $\varphi_h = 0$ on $]-1,1[\times]\frac{h}{2},1[$, and $||\nabla\varphi_h||_\infty \leq \frac{7}{h}$. Let us set $v_h := (1-\varphi_h)w_h$. We have that $v_h = 0$ in $\Omega' \setminus \overline{\Omega}$; moreover we have

$$\nabla v_h = (1-\varphi_h)\nabla w_h - \nabla\varphi_h w_h.$$

Since $\nabla\varphi_h w_h \to 0$ strongly in $L^2(\Omega'; \mathbb{R}^2)$, we have $\nabla v_h \to \nabla u$ strongly in $L^2(\Omega'; \mathbb{R}^N)$. Finally, for all A open subset of Ω' with $\mathcal{H}^1(\partial A \cap S_u) = 0$, we have

$$\lim_h \mathcal{H}^1(A \cap S_{v_h}) = \mathcal{H}^1(A \cap S_u).$$

In order to conclude the proof, let us apply Theorem 2.6 obtaining $\tilde{v}_h$ with polyhedral jumps in Ω such that $\tilde{v}_h \in W^{k,\infty}(\Omega' \setminus \overline{S_{\tilde{v}_h}})$, $||w_h - \tilde{v}_h||_{L^2(\Omega)} + ||\nabla w_h - \nabla\tilde{v}_h||_{L^2(\Omega;\mathbb{R}^2)} \leq h^2$ and $|\mathcal{H}^{N-1}(S_{w_h}) - \mathcal{H}^{N-1}(S_{\tilde{v}_h})| \leq h$. If we set $u_h := \varphi_h g + (1-\varphi_h)\tilde{v}_h$, we obtain the thesis. ☐

3. The discontinuous finite element approximation

In this section we construct a discrete approximation of quasi-static evolution of brittle fractures in linearly elastic bodies: the discretization is done both in space and time.

From now on we suppose that Ω is a polygonal open bounded subset of $\mathbb{R}^2$, and that $\partial_D\Omega \subseteq \partial\Omega$ is open in the relative topology. For all $\varepsilon > 0$, we fix a triangulation $\mathbf{R}_\varepsilon \in \mathcal{R}_\varepsilon(\Omega)$, and suppose that $\partial_D\Omega$ is composed of edges of $\mathbf{R}_\varepsilon$ for all ε; we indicate the family of these edges by $\mathbf{S}_\varepsilon$.

We consider the following discontinuous finite element space. We indicate by $\mathcal{A}_{\varepsilon,a}(\Omega)$ the set of all u such that there exists a triangulation $\mathbf{T}(u) \in \mathcal{T}_{\varepsilon,a}(\Omega)$ nested in $\mathbf{R}_\varepsilon$ with u affine on every $T \in \mathbf{T}(u)$. For every $u \in \mathcal{A}_{\varepsilon,a}(\Omega)$, we write $\|\nabla u\|$ for the L^2-norm of ∇u and we indicate by S_u the family of edges of $\mathbf{T}(u)$ inside Ω across which u is discontinuous. Notice that $u \in SBV(\Omega)$ and that the notation is consistent with the usual one employed in the theory of functions with bounded variation. Let us also denote by $\mathcal{AF}_\varepsilon(\Omega)$ the set of affine functions in Ω with respect to the triangulation $\mathbf{R}_\varepsilon$. Finally, given any $g \in \mathcal{AF}_\varepsilon(\Omega)$, for all $u \in \mathcal{A}_{\varepsilon,a}(\Omega)$ we set

$$S^g_D(u) := \{\zeta \in \mathbf{S}_\varepsilon \,:\, u \neq g \text{ on } \zeta\}, \tag{3.1}$$

that is $S^g_D(u)$ denotes the edges at which the boundary condition is not satisfied. Moreover we set

$$S^g(u) := S_u \cup S^g_D(u) \tag{3.2}$$

Let now consider $g \in W^{1,1}([0,1]; H^1(\Omega))$ with $g(t) \in \mathcal{AF}_\varepsilon(\Omega)$ for all $t \in [0,1]$. Let $\delta > 0$ and let N_δ be the largest integer such that $\delta(N_\delta - 1) < 1$; for $0 \le i \le N_\delta - 1$ we set $t_i^\delta := i\delta$, $t_{N_\delta}^\delta := 1$ and $g_i^\delta := g(t_i^\delta)$. The following proposition holds.

Proposition 3.1. *Let $\varepsilon > 0$, $a \in]0, \frac{1}{2}]$ and $\delta > 0$ be fixed. Then for all $i = 0, \dots, N_\delta$ there exists $u_{\varepsilon,a}^{\delta,i} \in \mathcal{A}_{\varepsilon,a}(\Omega)$ such that, setting*

$$\Gamma_{\varepsilon,a}^{\delta,i} := \bigcup_{r=0}^{i} S^{g_r^\delta}(u_{\varepsilon,a}^{\delta,r}),$$

the following hold:

(a) $\|u_{\varepsilon,a}^{\delta,i}\|_\infty \le \|g_i^\delta\|_\infty$;

(b) *for all $v \in \mathcal{A}_{\varepsilon,a}(\Omega)$ we have*

$$\|\nabla u_{\varepsilon,a}^{\delta,0}\|^2 + \mathcal{H}^1\left(S^{g_0^\delta}(u_{\varepsilon,a}^{\delta,0})\right) \le \|\nabla v\|^2 + \mathcal{H}^1\left(S^{g_0^\delta}(v)\right), \tag{3.3}$$

and

$$\|\nabla u_{\varepsilon,a}^{\delta,i}\|^2 \le \|\nabla v\|^2 + \mathcal{H}^1\left(S^{g_i^\delta}(v) \setminus \Gamma_{\varepsilon,a}^{\delta,i-1}\right). \tag{3.4}$$

Proof. The proof is carried out through a variational argument. Let $u_{\varepsilon,a}^{\delta,0}$ be a minimum of the following problem

$$\min\left\{\|\nabla u\|^2 + \mathcal{H}^1(S^{g_0^\delta}(u))\right\}. \tag{3.5}$$

We set $\Gamma_{\varepsilon,a}^{\delta,0} := S^{g_0^\delta}(u_{\varepsilon,a}^{\delta,0})$. Recursively, supposing to have constructed $u_{\varepsilon,a}^{\delta,i-1}$ and $\Gamma_{\varepsilon,a}^{\delta,i-1}$, let $u_{\varepsilon,a}^{\delta,i}$ be a minimum for

$$\min\left\{\|\nabla u\|^2 + \mathcal{H}^1\left(S^{g_i^\delta}(u) \setminus \Gamma_{\varepsilon,a}^{\delta,i-1}\right)\right\}. \tag{3.6}$$

We set $\Gamma_{\varepsilon,a}^{\delta,i} := S^{g_i^\delta}(u_{\varepsilon,a}^{\delta,i}) \cup \Gamma_{\varepsilon,a}^{\delta,i-1}$. We claim that problems (3.5) and (3.6) admit a solution $u_{\varepsilon,a}^{\delta,i}$ such that $\|u_{\varepsilon,a}^{\delta,i}\|_\infty \le \|g_i^\delta\|_\infty$ for all $i = 0, \dots, N_\delta$. We prove the claim for problem (3.6), the other case being similar. Let (u_n) be a minimizing sequence for problem (3.6): since g_i^δ is an admissible test function, we deduce that for n large

$$\|\nabla u_n\|^2 + \mathcal{H}^1\left(S^{g_i^\delta}(u_n) \setminus \Gamma_{\varepsilon,a}^{\delta,i-1}\right) \le \|\nabla g_i^\delta\|^2.$$

Moreover, we may modify u_n in the following way. If π denotes the projection in $\mathbb{R}$ over the interval $I := [-\|g_i^\delta\|_\infty, \|g_i^\delta\|_\infty]$, let $\tilde{u}_n \in \mathcal{A}_{\varepsilon,a}(\Omega)$ be defined on each $T \in \mathbf{T}(u_n)$ as the affine interpolation of the values $(\pi(u_n(x_1)),\ \pi(u_n(x_2)),\ \pi(u_n(x_3)))$, where x_1, x_2 and x_3 are the vertices of T. Note that by construction we have for all n

$$\|\tilde{u}_n\|_\infty \le \|g_i^\delta\|_\infty, \qquad \|\nabla \tilde{u}_n\| \le \|\nabla u_n\|, \qquad S^{g_i^\delta}(\tilde{u}_n) \subseteq S^{g_i^\delta}(u_n),$$

so that $(\tilde{u}_n)$ is a minimizing sequence for problem (3.6). We conclude that it is not restrictive to assume $\|u_n\|_\infty \le \|g_i^\delta\|_\infty$.

Since $\mathbf{T}(u_n) \in \mathcal{T}_{\varepsilon,a}(\Omega)$, we have that the number of elements of $\mathbf{T}(u_n)$ is uniformly bounded. Up to a subsequence, we may suppose that there exists an integer k such that $\mathbf{T}(u_n)$ has exactly k elements $T_n^1, \dots, T_n^k$. Using a diagonal argument we may suppose that, up to a further subsequence, there exists $\mathbf{T} = \{T^1, \dots, T^k\} \in \mathcal{T}_{\varepsilon,a}(\Omega)$ such that $T^i_n \to T^i$ in the Hausdorff metric for all $i = 1, \dots, k$. Let us consider $T^i \in \mathbf{T}$, and let $\tilde{T}^i$ be contained in the interior of T^i. For n large enough, $\tilde{T}^i$ is contained in the interior of T^i_n and $(u_n)_{|\tilde{T}^i}$ is affine with $\int_{\tilde{T}^i} |\nabla u_n|^2 \, dx \le C$ with C independent of n. We deduce that there exists a function u^i affine on $\tilde{T}^i$ such that up to a subsequence $u_n \to u$ uniformly on $\tilde{T}^i$. Since $\tilde{T}^i$ is arbitrary, it turns out that u^i is actually defined on T^i and

$$\int_{T^i} |\nabla u^i|^2 \, dx \le \liminf_n \int_{T^i_n} |\nabla u_n|^2 \, dx.$$

Let $u \in \mathcal{A}_{\varepsilon,a}(\Omega)$ such that $u = u^i$ on T^i for every $i = 1, \dots, k$: we have

$$\|\nabla u\|^2 \le \liminf_n \|\nabla u_n\|^2.$$

On the other hand, it is easy to see that $S^{g^\delta_i}(u)$ is contained in the Hausdorff limit of $S^{g^\delta_i}(u_n)$, and that

$$\mathcal{H}^1\left(S^{g^\delta_i}(u) \setminus \Gamma^{\delta,i-1}_{\varepsilon,a}\right) \le \liminf_n \mathcal{H}^1\left(S^{g^\delta_i}(u_n) \setminus \Gamma^{\delta,i-1}_{\varepsilon,a}\right).$$

We conclude that u is a minimum point for the problem (3.6) with $\|u\|_\infty \le \|g^\delta_i\|_\infty$. We have that point (a) is proved.

Concerning point (b), by construction we get (3.3); for $i \ge 1$ we have

$$\|\nabla u^{\delta,i}_{\varepsilon,a}\|^2 + \mathcal{H}^1\left(S^{g^\delta_i}(u^{\delta,i}_{\varepsilon,a}) \setminus \Gamma^{\delta,i-1}_{\varepsilon,a}\right) \le \|\nabla v\|^2 + \mathcal{H}^1\left(S^{g^\delta_i}(v) \setminus \Gamma^{\delta,i-1}_{\varepsilon,a}\right)$$

for all $v \in \mathcal{A}_{\varepsilon,a}(\Omega)$, so that

$$\|\nabla u^{\delta,i}_{\varepsilon,a}\|^2 \le \|\nabla v\|^2 + \mathcal{H}^1\left(S^{g^\delta_i}(v) \setminus \Gamma^{\delta,i-1}_{\varepsilon,a}\right),$$

and this proves point (b). $\square$

Remark 3.2. For technical reasons due to the asymptotic analysis of the discrete evolution $u^{\delta,i}_{\varepsilon,a}$ when $\delta \to 0$, $\varepsilon \to 0$ and $a \to 0$, we define $u^{\delta,i}_{\varepsilon,a}$ from $u^{\delta,i-1}_{\varepsilon,a}$ through problem (3.6) without requiring that the adaptive vertices determining $\Gamma^{\delta,i-1}_{\varepsilon,a}$ remain fixed. We just penalize their possible changes if they are used to create new fracture: in fact in this case, the surface energy increases at each change of a quantity at least of order $a\varepsilon$. As a consequence, during the step by step minimization, it could happen that some triangles $T \in \mathcal{T}_{\varepsilon,a}(\Omega)$ contain the fracture $\Gamma^{\delta,i}_{\varepsilon,a}$ in their interior. This is in contrast with the interpretation of the triangles as elementary blocks for the elasticity problem, but being this situation penalized in the minimization process, we expect that it occurs rarely.

The following estimate is essential for the study of asymptotic behavior of the discrete evolution.

Proposition 3.3. *If* $(u^{\delta,i}_{\varepsilon,a}, \Gamma^{\delta,i}_{\varepsilon,a})$ *for* $i = 0, \dots, N_\delta$ *satisfies condition* (b) *of Proposition 3.1, setting* $\mathcal{E}^{\delta,i}_{\varepsilon,a} := \|\nabla u^{\delta,i}_{\varepsilon,a}\|^2 + \mathcal{H}^1\left(\Gamma^{\delta,i}_{\varepsilon,a}\right)$*, we have for* $0 \le j \le i \le N_\delta$

$$\mathcal{E}^{\delta,i}_{\varepsilon,a} \le \mathcal{E}^{\delta,j}_{\varepsilon,a} + 2\sum_{r=j}^{i-1} \int_{t^\delta_r}^{t^\delta_{r+1}} \int_\Omega \nabla u^{\delta,r}_{\varepsilon,a} \nabla \dot{g}(\tau) \, dx \, d\tau + o^\delta, \tag{3.7}$$

where

$$o^\delta := \left[\max_{r=0,\dots,N_\delta-1} \int_{t_r^\delta}^{t_{r+1}^\delta} \|\dot g(\tau)\|_{H^1(\Omega)}\,d\tau \right] \int_0^1 \|\dot g(\tau)\|_{H^1(\Omega)}\,d\tau. \tag{3.8}$$

Proof. For all $0 \le j \le N_\delta - 1$, by construction of $u_{\varepsilon,a}^{\delta,j+1}$ we have that

$$\|\nabla u_{\varepsilon,a}^{\delta,j+1}\|^2 + \mathcal{H}^1\left(S^{g_{j+1}^\delta}(u_{\varepsilon,a}^{\delta,j+1}) \setminus \Gamma_{\varepsilon,a}^{\delta,j} \right) \le \|\nabla u_{\varepsilon,a}^{\delta,j} + \nabla(g_{j+1}^\delta - g_j^\delta)\|^2 =$$
$$= \|\nabla u_{\varepsilon,a}^{\delta,j}\|^2 + 2\int_\Omega \nabla u_{\varepsilon,a}^{\delta,j} \nabla(g_{j+1}^\delta - g_j^\delta)\,dx + ||\nabla(g_{j+1}^\delta - g_j^\delta)||^2.$$

Notice that

$$\nabla(g_{j+1}^\delta - g_j^\delta) = \int_{t_j^\delta}^{t_{j+1}^\delta} \nabla \dot g(\tau)\,d\tau,$$

so that

$$\|\nabla u_{\varepsilon,a}^{\delta,j+1}\|^2 + \mathcal{H}^1\left(S^{g_{j+1}^\delta}(u_{\varepsilon,a}^{\delta,j+1}) \setminus \Gamma_{\varepsilon,a}^{\delta,j} \right) \le \tag{3.9}$$
$$\le \|\nabla u_{\varepsilon,a}^{\delta,j}\|^2 + 2\int_{t_j^\delta}^{t_{j+1}^\delta} \int_\Omega \nabla u_{\varepsilon,a}^{\delta,j} \nabla \dot g(\tau)\,dx\,d\tau + e(\delta) \int_{t_j^\delta}^{t_{j+1}^\delta} \|\dot g(\tau)\|_{H^1(\Omega)}\,d\tau,$$

where

$$e(\delta) := \max_{r=0,\dots,N_\delta-1} \int_{t_r^\delta}^{t_{r+1}^\delta} \|\dot g(\tau)\|_{H^1(\Omega)}\,d\tau.$$

From (3.9), we obtain that for all $0 \le j \le i \le N_\delta$

$$||\nabla u_{\varepsilon,a}^{\delta,i}||^2 + \mathcal{H}^1(\Gamma_{\varepsilon,a}^{\delta,i}) \le ||\nabla u_{\varepsilon,a}^{\delta,j}||^2 + \mathcal{H}^1(\Gamma_{\varepsilon,a}^{\delta,j}) +$$
$$+ 2\sum_{r=j}^{i-1} \int_{t_r^\delta}^{t_{r+1}^\delta} \int_\Omega \nabla u_{\varepsilon,a}^{\delta,r} \nabla \dot g(\tau)\,dx\,d\tau + e(\delta) \int_{t_j^\delta}^{t_i^\delta} ||\dot g(\tau)||_{H^1(\Omega)}\,d\tau,$$

and so the proof of point (c) is complete choosing

$$o^\delta := e(\delta) \int_0^1 \|\dot g(\tau)\|_{H^1(\Omega)}\,d\tau.$$

□

4. THE CONVERGENCE RESULT

This section is devoted to the proof of Theorem 1.1. As in Section 3, let Ω be a polygonal open bounded subset of $\mathbb{R}^2$, and let $\partial_D\Omega \subseteq \partial\Omega$ be open in the relative topology. For all $\varepsilon > 0$, let $\mathbf{R}_\varepsilon \in \mathcal{R}_\varepsilon(\Omega)$ be a regular triangulation of Ω such that $\partial_D\Omega$ is composed of edges of $\mathbf{R}_\varepsilon$. As in the previous section, let $\mathcal{AF}_\varepsilon(\Omega)$ be the family of continuous piecewise affine functions with respect to $\mathbf{R}_\varepsilon$, and let $\mathcal{A}_{\varepsilon,a}(\Omega)$ be the family of functions which are affine on the triangles of some triangulation $\mathbf{T} \in \mathcal{T}_{\varepsilon,a}(\Omega)$ nested in $\mathbf{R}_\varepsilon$ and can jump across the edges of $\mathbf{T}$.

In the following, it will be useful to treat points at which the boundary condition is violated (see (3.1)) as internal jumps. Thus we consider Ω_D polygonal open bounded subset of $\mathbb{R}^2$ such that $\Omega_D \cap \Omega = \emptyset$ and $\partial\Omega \cap \partial\Omega_D = \partial_D\Omega$ up to a finite number of points; we set $\Omega' := \Omega \cup \Omega_D \cup \partial_D\Omega$. Given $u \in \mathcal{A}_{\varepsilon,a}(\Omega)$ and $g \in \mathcal{AF}_\varepsilon(\Omega)$, we may extend g to a function of

$H^1(\Omega')$ and u to a function $\tilde{u} \in SBV(\Omega')$ setting $\tilde{u} = g$ on Ω_D. In this way, recalling (3.2), we have

$$S^g(u) = S_{\tilde{u}},$$

so that the violation of the boundary condition of u can be read in the set of jumps of $\tilde{u}$. Analogously, given $u \in SBV(\Omega)$ and $g \in H^1(\Omega)$, we set

$$S^g(u) := S_u \cup \{x \in \partial_D\Omega \, : \, \gamma(u)(x) \neq \gamma(g)(x)\} \tag{4.1}$$

where γ denotes the trace operator on $\partial\Omega$. We may assume $g \in H^1(\Omega')$ using an extension operator. We can then consider $\tilde{u} \in SBV(\Omega')$ such that $\tilde{u} = u$ on Ω, and $\tilde{u} = g$ on Ω_D. In this way we have

$$S^g(u) = S_{\tilde{u}} \quad \text{up to a set of } \mathcal{H}^1\text{-measure } 0.$$

Let us consider $g \in W^{1,1}([0,1], H^1(\Omega))$ such that $\|g(t)\|_\infty \leq C$ for all $t \in [0,1]$ and let $g_\varepsilon \in W^{1,1}([0,1], H^1(\Omega))$ be such that $g_\varepsilon(t) \in \mathcal{AF}_\varepsilon(\Omega)$ for all $t \in [0,1]$,

$$\|g_\varepsilon(t)\|_\infty \leq C \tag{4.2}$$

for all $t \in [0,1]$, and for $\varepsilon \to 0$

$$g_\varepsilon \to g \quad \text{strongly in } W^{1,1}([0,1], H^1(\Omega)). \tag{4.3}$$

We indicate by $\{u^{\delta,i}_{\varepsilon,a}, \, i = 0, \dots, N_\delta\}$ the discrete evolution relative to the boundary data g_ε given by Proposition 3.1, and we denote by $\mathcal{E}^{\delta,i}_{\varepsilon,a}$ its total energy as in Proposition 3.3.

We assume that $g(\cdot)$ and $g_\varepsilon(\cdot)$ are defined in $H^1(\Omega')$ (we still denote these extensions by $g(\cdot)$ and $g_h(\cdot)$), in such a way that (4.2) and (4.3) hold in Ω'. Let us moreover set $g^\delta_\varepsilon(t) := g_\varepsilon(t^\delta_i)$ for all $t^\delta_i \leq t < t^\delta_{i+1}$ with $i = 0, \dots, N_\delta - 1$ and $g^\delta_\varepsilon(1) := g_\varepsilon(1)$.

Let us make the following piecewise constant interpolation in time:

$$u^\delta_{\varepsilon,a}(t) := u^{\delta,i}_{\varepsilon,a} \quad \text{for } t^\delta_i \leq t < t^\delta_{i+1} \quad i = 0, \dots, N_\delta - 1,$$

and $u^\delta_{\varepsilon,a}(1) := u^{\delta,N_\delta}_{\varepsilon,a}$. For all $t \in [0,1]$ we define the *discrete fracture* at time t as

$$\Gamma^\delta_{\varepsilon,a}(t) := \bigcup_{s \leq t} S^{g^\delta_\varepsilon(s)}(u^\delta_{\varepsilon,a}(s)),$$

and the *discrete total energy* at time t as

$$\mathcal{E}^\delta_{\varepsilon,a}(t) := \|\nabla u^\delta_{\varepsilon,a}(t)\|^2 + \mathcal{H}^1\left(\Gamma^\delta_{\varepsilon,a}(t)\right).$$

We have for all $t \in [0,1]$

$$\|u^\delta_{\varepsilon,a}(t)\|_\infty \leq \|g^\delta_\varepsilon(t)\|_\infty. \tag{4.4}$$

Moreover for all $v \in \mathcal{A}_{\varepsilon,a}(\Omega)$ we have

$$\|\nabla u^\delta_{\varepsilon,a}(0)\|^2 + \mathcal{H}^1\left(S^{g^\delta_\varepsilon(0)}(u^\delta_{\varepsilon,a}(0))\right) \leq \|\nabla v\|^2 + \mathcal{H}^1\left(S^{g^\delta_\varepsilon(0)}(v)\right), \tag{4.5}$$

and for all $t \in]0,1]$ and for all $v \in \mathcal{A}_{\varepsilon,a}(\Omega)$

$$\|\nabla u^\delta_{\varepsilon,a}(t)\|^2 \leq \|\nabla v\|^2 + \mathcal{H}^1\left(S^{g^\delta_\varepsilon(t)}(v) \setminus \Gamma^\delta_{\varepsilon,a}(t)\right). \tag{4.6}$$

Finally for all $0 \leq s \leq t \leq 1$ we have

$$\mathcal{E}^\delta_{\varepsilon,a}(t) \leq \mathcal{E}^\delta_{\varepsilon,a}(s) + 2\int_{s^\delta_i}^{t^\delta_i} \int_\Omega \nabla u^\delta_{\varepsilon,a}(\tau) \nabla \dot{g}_\varepsilon(\tau) \, dx \, d\tau + o^\delta_\varepsilon, \tag{4.7}$$

where $t_i^\delta \leq t < t_{i+1}^\delta$, $s_i^\delta \leq s < s_{i+1}^\delta$ and

$$o_\varepsilon^\delta := \left[\max_{r=0,\dots,N_\delta-1} \int_{t_r^\delta}^{t_{r+1}^\delta} \|\dot{g}_\varepsilon(\tau)\|_{H^1(\Omega)}\, d\tau \right] \int_0^1 \|\dot{g}_\varepsilon(\tau)\|_{H^1(\Omega)}. \tag{4.8}$$

For $s = 0$ we obtain the following estimate from above for the discrete total energy

$$\mathcal{E}_{\varepsilon,a}^\delta(t) \leq \mathcal{E}_{\varepsilon,a}^\delta(0) + 2\int_0^{t_i^\delta} \int_\Omega \nabla u_{\varepsilon,a}^\delta(\tau) \nabla \dot{g}_\varepsilon(\tau)\, dx\, d\tau + o_\varepsilon^\delta, \tag{4.9}$$

where $t_i^\delta \leq t < t_{i+1}^\delta$.

We study the behavior of the evolution $\{t \to u_{\varepsilon,a}^\delta(t),\, t \in [0,1]\}$ varying the parameters in the following way. We let firstly $\varepsilon \to 0$ and $\delta \to 0$ obtaining an evolution $\{t \to u_a(t),\, t \in [0,1]\}$ relative to the boundary data g with the minimality property (4.17); then we let $a \to 0$ obtaining a quasi-static evolution of brittle fractures $\{t \to u(t)\,, t \in [0,1]\}$ relative to the boundary data g. Finally, by a diagonal argument we deal with (δ, ε, a) at the same time.

In order to develop this program, we need some compactness, and so we derive a bound for the total energy $\mathcal{E}_{\varepsilon,a}^\delta$. By (3.4), we have that for all $t \in [0,1]$

$$\|\nabla u_{\varepsilon,a}^\delta(t)\| \leq \|\nabla g_\varepsilon^\delta(t)\| \leq \tilde{C}$$

with $\tilde{C}$ independent of δ, ε and t. We deduce for all $t \in [0,1]$

$$\mathcal{E}_{\varepsilon,a}^\delta(t) \leq \mathcal{E}_{\varepsilon,a}^\delta(0) + 2\tilde{C}^2 + o_\varepsilon^\delta$$

Notice that $\mathcal{E}_{\varepsilon,a}^\delta(0)$ is uniformly bounded as δ, ε vary. Moreover, by (4.4) and since $\|g_\varepsilon(t)\|_\infty \leq C$ for all $t \in [0,1]$, we have that $u_{\varepsilon,a}^\delta(t)$ is uniformly bounded in $L^\infty(\Omega)$ independently of δ, ε and a. Taking into account (4.3), we conclude that there exists C' independent of δ, ε, a such that for all $t \in [0,1]$

$$\mathcal{E}_{\varepsilon,a}^\delta(t) + \|u_{\varepsilon,a}^\delta(t)\|_\infty \leq C'. \tag{4.10}$$

Formula (4.10) gives the desired compactness in order to perform the asymptotic analysis of the discrete evolution.

Let now consider $\delta_n \to 0$ and $\varepsilon_n \to 0$: by (4.3) we have

$$o_{\varepsilon_n}^{\delta_n} \to 0, \tag{4.11}$$

where $o_{\varepsilon_n}^{\delta_n}$ is defined in (4.8). By Helly's theorem on monotone functions, we may suppose that there exists an increasing function λ_a such that (up to a subsequence) for all $t \in [0,1]$

$$\lambda_{n,a}(t) := \mathcal{H}^1\left(\bigcup_{s \leq t} S^{g_{\varepsilon_n}^{\delta_n}(s)}(u_{\varepsilon_n,a}^{\delta_n}(s)) \right) \to \lambda_a(t). \tag{4.12}$$

Let us fix $D \subseteq [0,1]$ countable and dense with $0 \in D$.

Lemma 4.1. *For all $t \in D$ there exists $u_a(t) \in SBV(\Omega)$ such that up to a subsequence independent of t*

$$u_{\varepsilon_n,a}^{\delta_n}(t) \to u_a(t) \quad \text{in } SBV(\Omega).$$

Moreover for all $t \in D$ we have

$$\|\nabla u_a(t)\|^2 + \mathcal{H}^1\big(S^{g(t)}(u_a(t))\big) + \|u_a(t)\|_\infty \leq C'. \tag{4.13}$$

Proof. Let us consider $t \in D$. By (4.10), we can apply Ambrosio's Compactness Theorem 2.1 obtaining $u \in SBV(\Omega)$ such that, up to a subsequence, $u^{\delta_n}_{\varepsilon_n,a}(t) \to u$ in $SBV(\Omega)$. Let us set $u_a(t) := u$. Using a diagonal argument, we deduce that there exists a subsequence of $(\delta_n, \varepsilon_n)$ (which we still denote by $(\delta_n, \varepsilon_n)$) such that $u^{\delta_n}_{\varepsilon_n,a}(t) \to u_a(t)$ in $SBV(\Omega)$ for all $t \in D$. In order to obtain inequality (4.13), we extend $u^{\delta_n}_{\varepsilon_n,a}(t)$ and $u_a(t)$ to Ω' setting $u^{\delta_n}_{\varepsilon_n,a}(t) := g^{\delta_n}_{\varepsilon_n}(t)$ and $u_a(t) := g(t)$ on Ω_D; since $g^{\delta_n}_{\varepsilon_n}(t) \to g(t)$ on Ω_D strongly in $H^1(\Omega_D)$, we have that $u^{\delta_n}_{\varepsilon_n,a}(t) \to u_a(t)$ in $SBV(\Omega')$, so that we can apply Ambrosio's Theorem, and derive (4.13) from (4.10). ☐

The following result is essential for the sequel: its proof is postponed to Section 5.

Proposition 4.2. *Let $t \in D$. For all $v \in SBV(\Omega)$ we have*

$$\|\nabla u_a(t)\|^2 \le \|\nabla v\|^2 + \mu(a)\mathcal{H}^1(S^{g(t)}(v) \setminus \bigcup_{s \le t, s \in D} S^{g(s)}(u_a(s))), \tag{4.14}$$

where $\mu :]0, \frac{1}{2}[\to]0, +\infty[$ is such that $\lim_{a \to 0} \mu(a) = 1$. Moreover, $\nabla u^{\delta_n}_{\varepsilon_n,a}(t) \to \nabla u_a(t)$ strongly in $L^2(\Omega; \mathbb{R}^2)$.

We now extend the evolution $\{t \to u_a(t) : t \in D\}$ to the entire interval $[0, 1]$. Let us set for all $t \in [0, 1]$

$$\Gamma_a(t) := \bigcup_{s \le t, s \in D} S^{g(s)}(u_a(s)).$$

Lemma 4.3. *For every $t \in [0, 1]$ there exists $u_a(t) \in SBV(\Omega)$ such that the following hold:*

(a) *for all $t \in [0, 1]$*

$$S^{g(t)}(u_a(t)) \subseteq \Gamma_a(t) \text{ up to a set of } \mathcal{H}^1\text{-measure } 0, \tag{4.15}$$

and

$$\|\nabla u_a(t)\|^2 + \mathcal{H}^1\big(S^{g(t)}(u_a(t))\big) + \|u_a(t)\|_\infty \le C'; \tag{4.16}$$

(b) *for all $v \in SBV(\Omega)$*

$$\|\nabla u_a(t)\|^2 \le \|\nabla v\|^2 + \mu(a)\mathcal{H}^1\left(S^{g(t)}(v) \setminus \Gamma_a(t)\right); \tag{4.17}$$

(c) *∇u_a is left continuous in $[0, 1] \setminus D$ with respect to the strong topology of $L^2(\Omega; \mathbb{R}^2)$;*

(d) *for all $t \in [0, 1] \setminus \mathcal{N}_a$ we have that*

$$\nabla u^{\delta_n}_{\varepsilon_n,a}(t) \to \nabla u_a(t) \quad \text{strongly in } L^2(\Omega, \mathbb{R}^2),$$

where $\mathcal{N}_a$ is the set of discontinuities of the function λ_a defined in (4.12).

Proof. Let $t \in [0, 1] \setminus D$ and let $t_n \in D$ with $t_n \nearrow t$. By (4.13), we can apply Ambrosio's Theorem to the sequence $(u_a(t_n))$ obtaining $u \in SBV(\Omega)$ such that, up to a subsequence, $u_a(t_n) \to u$ in $SBV(\Omega)$. Let us set $u_a(t) := u$. Let us extend $u_a(t_n)$ and $u_a(t)$ to Ω' setting $u_a(t_n) := g(t_n)$ and $u_a(t) := g(t)$ on Ω_D: we have $u_a(t_n) \to u_a(t)$ in $SBV(\Omega')$. Since $\mathcal{H}^1 \llcorner S_{u_a(t_n)} \le \mathcal{H}^1 \llcorner \Gamma_a(t)$ for all n, as a consequence of Ambrosio's Theorem, we deduce that

$\mathcal{H}^1 \llcorner S_{u_a(t)} \leq \mathcal{H}^1 \llcorner \Gamma_a(t)$. This means $\mathcal{H}^1 \llcorner S^{g(t)}(u_a(t)) \leq \mathcal{H}^1 \llcorner \Gamma_a(t)$, so that (4.15) holds. Moreover, for all $v \in SBV(\Omega)$, by (4.14) we may write

$$\begin{aligned} \|\nabla u_a(t_n)\|^2 &\leq \|\nabla v - \nabla g(t) + \nabla g(t_n)\|^2 + \mu(a)\mathcal{H}^1\left(S^{g(t)}(v) \setminus \Gamma_a(t_n)\right) \leq \\ &\leq \|\nabla v - \nabla g(t) + \nabla g(t_n)\|^2 + \mu(a)\mathcal{H}^1\left(S^{g(t)}(v) \setminus \Gamma_a(t)\right) + \mu(a)\mathcal{H}^1\left(\Gamma_a(t) \setminus \Gamma_a(t_n)\right), \end{aligned} \tag{4.18}$$

so that, since by definition of $\Gamma_a(t)$ we have $\mathcal{H}^1(\Gamma_a(t) \setminus \Gamma_a(t_n)) \to 0$, we obtain that (4.17) holds; choosing $v = u_a(t)$ and taking the limsup in (4.18), we obtain that

$$\limsup_n \|\nabla u_a(t_n)\|^2 \leq \|\nabla u_a(t)\|^2,$$

and so the convergence $\nabla u_a(t_n) \to \nabla u_a(t)$ is strong in $L^2(\Omega, \mathbb{R}^2)$. Notice that $\nabla u_a(t)$ is uniquely determined by (4.15) and (4.17) since the gradient of the solutions of the minimum problem

$$\min\left\{\|\nabla u\|^2 \,:\, S^{g(t)}(u) \subseteq \Gamma_a(t) \text{ up to a set of } \mathcal{H}^1\text{-measure } 0\right\}$$

is unique by the strict convexity of the functional: we conclude that $\nabla u_a(t)$ is well defined. The same arguments prove that ∇u_a is left continuous at all $t \in [0,1] \setminus D$. Finally (4.16) is a direct consequence of (4.13) and of Ambrosio's Theorem, and so points (a), (b), (c) are proved.

Let us come to point (d). Let us consider $u^{\delta_n}_{\varepsilon_n,a}(t)$ with $t \notin \mathcal{N}_a$; we may suppose that $t \notin D$, since otherwise the result has already been established. By Proposition 4.2 with $D' := D \cup \{t\}$ in place of D, we have that, up to a subsequence, $u^{\delta_n}_{\varepsilon_n,a}(t) \to u$ in $SBV(\Omega)$ such that

$$\|\nabla u\|^2 \leq \|\nabla v\|^2 + \mu(a)\mathcal{H}^1\left(S^{g(t)}(v) \setminus (\Gamma_a(t) \cup S^{g(t)}(u))\right)$$

for all $v \in SBV(\Omega)$ and $\nabla u^{\delta_n}_{\varepsilon_n,a}(t) \to \nabla u$ strongly in $L^2(\Omega; \mathbb{R}^2)$. Let $s < t$ with $s \in D$; by the minimality of $u^{\delta_n}_{\varepsilon_n,a}(s)$ and by (4.10) we have

$$\begin{aligned} \|\nabla u^{\delta_n}_{\varepsilon_n,a}(s)\|^2 &\leq \|\nabla u^{\delta_n}_{\varepsilon_n,a}(t) - \nabla g^{\delta_n}_{\varepsilon_n}(t) + \nabla g^{\delta_n}_{\varepsilon_n}(s)\|^2 + \lambda_{n,a}(t) - \lambda_{n,a}(s) \leq \\ &\leq \|\nabla u^{\delta_n}_{\varepsilon_n,a}(t)\|^2 + 2\sqrt{C'}\|\nabla g^{\delta_n}_{\varepsilon_n}(t) - \nabla g^{\delta_n}_{\varepsilon_n}(s)\| + \\ &\qquad + \|\nabla g^{\delta_n}_{\varepsilon_n}(t) - \nabla g^{\delta_n}_{\varepsilon_n}(s)\|^2 + \lambda_{n,a}(t) - \lambda_{n,a}(s). \end{aligned}$$

Passing to the limit for $n \to +\infty$, recalling that $g^{\delta_n}_{\varepsilon_n}(\tau) \to g(\tau)$ strongly in $H^1(\Omega)$ for all $\tau \in [0,1]$, we deduce

$$\|\nabla u_a(s)\|^2 \leq \|\nabla u\|^2 + 2\sqrt{C'}\|\nabla g(t) - \nabla g(s)\| + \|\nabla g(t) - \nabla g(s)\|^2 + \lambda_a(t) - \lambda_a(s),$$

so that, since t is a point of continuity for λ_a, ∇u_a is left continuous at t, and g is absolutely continuous, we get for $s \to t$

$$\|\nabla u_a(t)\|^2 \leq \|\nabla u\|^2.$$

We conclude that $u_a(t)$ is a solution of

$$\min\{\|\nabla v\|^2 \,:\, S^{g(t)}(v) \subseteq \Gamma_a(t) \cup S^{g(t)}(u) \text{ up to a set of } \mathcal{H}^1\text{-measure } 0\},$$

so that $\nabla u = \nabla u_a(t)$ by uniqueness of the gradient of the solution. We deduce that $\nabla u^{\delta_n}_{\varepsilon_n,a}(t) \to \nabla u_a(t)$ strongly in $L^2(\Omega; \mathbb{R}^2)$, and so the proof is complete. □

We can now let $a \to 0$.

Lemma 4.4. *There exists $a_n \to 0$ such that, for all $t \in D$, $u_{a_n}(t) \to u(t)$ in $SBV(\Omega)$ for some $u(t) \in SBV(\Omega)$ such that for all $v \in SBV(\Omega)$ we have*

$$\|\nabla u(t)\|^2 \leq \|\nabla v\|^2 + \mathcal{H}^1(S^{g(t)}(v) \setminus \bigcup_{s \leq t, s \in D} S^{g(s)}(u(s))). \tag{4.19}$$

Moreover, $\nabla u_{a_n}(t) \to \nabla u(t)$ strongly in $L^2(\Omega;\mathbb{R}^2)$ and

$$\|\nabla u(t)\|^2 + \mathcal{H}^1(S^{g(t)}(u(t))) + \|u(t)\|_\infty \leq C'. \tag{4.20}$$

Proof. By (4.16), applying Ambrosio's Theorem to the extensions of $u_a(t)$ to Ω' by setting $u_a(t) := g(t)$ on Ω_D, and using a diagonal argument, we find a sequence $a_n \to 0$ such that, for all $t \in D$, $u_{a_n}(t) \to u(t)$ in $SBV(\Omega)$ for some $u(t) \in SBV(\Omega)$ such that (4.20) holds.

We now prove that $u(t)$ satisfies property (4.19). Let $v \in SBV(\Omega)$. Let us fix $t_1 \leq t_2 \leq \ldots \leq t_k = t$ with $t_i \in D$. We extend v and $u_{a_n}(t_i)$ to Ω' setting $v := g(t)$ and $u_{a_n}(t_i) := g(t_i)$ on Ω_D respectively. Since $u_{a_n}(t_i) \to u(t_i)$ in $SBV(\Omega')$ for all $i = 1, \ldots, k$, by Theorem 2.3 there exists $v_n \in SBV(\Omega')$ with $v_n = g(t)$ on Ω_D such that $\nabla v_n \to \nabla v$ strongly in $L^2(\Omega';\mathbb{R}^2)$ and

$$\limsup_n \mathcal{H}^1\left(S_{v_n} \setminus \bigcup_{i=1}^k S_{u_{a_n}(t_i)}\right) \leq \mathcal{H}^1\left(S_v \setminus \bigcup_{i=1}^k S_{u(t_i)}\right). \tag{4.21}$$

By (4.14) we obtain

$$\|\nabla u_{a_n}(t)\|^2 \leq \|\nabla v_n\|^2 + \mu(a_n)\mathcal{H}^1\left(S_{v_n} \setminus \bigcup_{i=1}^k S_{u_{a_n}(t_i)}\right), \tag{4.22}$$

so that passing to the limit for $n \to +\infty$ and recalling that $\mu(a) \to 1$ as $a \to 0$, we obtain

$$\|\nabla u(t)\|^2 \leq \|\nabla v\|^2 + \mathcal{H}^1\left(S_v \setminus \bigcup_{i=1}^k S_{u(t_i)}\right).$$

Thus we get

$$\|\nabla u(t)\|^2 \leq \|\nabla v\|^2 + \mathcal{H}^1\left(S^{g(t)}(v) \setminus \bigcup_{i=1}^k S^{g(t_i)}(u(t_i))\right).$$

Since $t_1, \ldots, t_k$ are arbitrary, we obtain (4.19). Choosing $v = u(t)$, taking the limsup in (4.22) and using (4.21), we obtain $\nabla u_{a_n}(t) \to \nabla u(t)$ strongly in $L^2(\Omega;\mathbb{R}^2)$. ☐

In order to deal with δ, ε and a at the same time, we need the following lemma.

Lemma 4.5. *Let $\{u(t) : t \in D\}$ be as in Lemma 4.4. There exist $\delta_n \to 0$, $\varepsilon_n \to 0$, and $a_n \to 0$ such for all $t \in D$ we have*

$$u^{\delta_n}_{\varepsilon_n, a_n}(t) \to u(t) \quad \text{in } SBV(\Omega)\ .$$

Moreover, for all n there exists $\mathcal{B}_n \subseteq [0,1]$ with $|\mathcal{B}_n| < 2^{-n}$ such that for all $t \in [0,1] \setminus \mathcal{B}_n$

$$\|\nabla u^{\delta_n}_{\varepsilon_n, a_n}(t) - \nabla u_{a_n}(t)\| \leq \frac{1}{n}. \tag{4.23}$$

Finally, we have that for all $v \in SBV(\Omega)$

$$\|\nabla u(0)\|^2 + \mathcal{H}^1\left(S^{g(0)}(u(0))\right) \leq \|\nabla v\|^2 + \mathcal{H}^1\left(S^{g(0)}(v)\right) \tag{4.24}$$

and

$$\mathcal{E}^{\delta_n}_{\varepsilon_n,a_n}(0) \to \|\nabla u(0)\|^2 + \mathcal{H}^1\left(S^{g(0)}(u(0))\right). \tag{4.25}$$

Proof. Let (a_n) be the sequence determined by Lemma 4.4. By Lemma 4.1, for all n there exists $(\delta^n_m, \varepsilon^n_m)$ such that for all $t \in D$ and $m \to +\infty$ we have

$$u^{\delta^n_m}_{\varepsilon^n_m,a_n}(t) \to u_{a_n}(t) \quad \text{in } SBV(\Omega),$$

and

$$\nabla u^{\delta^n_m}_{\varepsilon^n_m,a_n}(t) \to \nabla u_{a_n}(t) \quad \text{strongly in } L^2(\Omega;\mathbb{R}^2).$$

Moreover by Lemma 4.3 we have that $\nabla u^{\delta^n_m}_{\varepsilon^n_m,a_n} \to \nabla u_{a_n}$ quasi-uniformly on $[0,1]$ as $m \to +\infty$. Let $\mathcal{B}_n \subseteq [0,1]$ with $|\mathcal{B}_n| < 2^{-n}$ such that $\nabla u^{\delta^n_m}_{\varepsilon^n_m,a_n} \to \nabla u_{a_n}$ uniformly on $[0,1] \setminus \mathcal{B}_n$ as $m \to +\infty$. We now perform the following diagonal argument. Let $D = \{t_n,\, n \geq 1\}$. Choose m_1 such that

$$\|\nabla u^{\delta^1_{m_1}}_{\varepsilon^1_{m_1},a_1}(t_1) - \nabla u_{a_1}(t_1)\| + \|u^{\delta^1_{m_1}}_{\varepsilon^1_{m_1},a_1}(t_1) - u_{a_1}(t_1)\| \leq 1,$$

and

$$\|\nabla u^{\delta^1_{m_1}}_{\varepsilon^1_{m_1},a_1}(t) - \nabla u_{a_1}(t)\| \leq 1 \quad \text{for all } t \in [0,1] \setminus \mathcal{B}_1 .$$

Let m_n be such that

$$\|\nabla u^{\delta^n_{m_n}}_{\varepsilon^n_{m_n},a_n}(t_j) - \nabla u_{a_n}(t_j)\| + \|u^{\delta^n_{m_n}}_{\varepsilon^n_{m_n},a_n}(t_j) - u_{a_n}(t_j)\| \leq \frac{1}{n} \quad \text{for all } j = 1, \dots, n$$

and

$$\|\nabla u^{\delta^n_{m_n}}_{\varepsilon^n_{m_n},a_n}(t) - \nabla u_{a_n}(t)\| \leq \frac{1}{n} \quad \text{for all } t \in [0,1] \setminus \mathcal{B}_n .$$

We may suppose that $\delta^n_{m_n} \to 0$, $\varepsilon^n_{m_n} \to 0$. Then $(\delta^n_{m_n}, \varepsilon^n_{m_n}, a_n)$ is the sequence which satisfies the thesis. In fact by construction and taking into account (4.10), for all $t \in D$ we have $u^{\delta^n_{m_n}}_{\varepsilon^n_{m_n},a_n}(t) \to u(t)$ in $SBV(\Omega)$; moreover the set $\mathcal{B}_n$ satisfies (4.23). Notice that $u^{\delta^n_{m_n}}_{\varepsilon^n_{m_n},a^n_{m_n}}(0)$ satisfies (4.5) and so (4.24) and (4.25) follow by the Γ-convergence result of [12]. $\square$

Let $(\delta_n, \varepsilon_n, a_n)$ be the sequence determined by Lemma 4.5. For all $t \in [0,1]$ let us set

$$\lambda_n(t) := \mathcal{H}^1\left(\Gamma^{\delta_n}_{\varepsilon_n,a_n}(t)\right).$$

By Helly's theorem, we may suppose that there exist two increasing functions λ and η such that up to a subsequence

$$\lambda_n \to \lambda \quad \text{pointwise in } [0,1],$$

and

$$\lambda_{a_n} \to \eta \quad \text{pointwise in } [0,1], \tag{4.26}$$

where λ_{a_n} is defined as in (4.12). We now extend the evolution $\{t \to u(t) \,:\, t \in D\}$ to the entire interval $[0,1]$. Let us set for all $t \in [0,1]$

$$\Gamma(t) := \bigcup_{s \leq t, s \in D} S^{g(s)}(u(s)),$$

and let $\mathcal{N}$ be the set of discontinuities of $\mathcal{H}^1(\Gamma(\cdot))$. Notice that for all $t \in [0,1]$

$$\mathcal{H}^1(\Gamma(t)) \leq \lambda(t). \tag{4.27}$$

In fact if $t \in D$, let $t_1 \leq t_2 \leq \ldots \leq t_k = t$ with $t_i \in D$, consider $w_n \in SBV(\Omega'; \mathbb{R}^k)$ defined as

$$w_n(x) := (u^{\delta_n}_{\varepsilon_n, a_n}(t_1)(x), \ldots, u^{\delta_n}_{\varepsilon_n, a_n}(t_k)(x)),$$

where we assume that $u^{\delta_n}_{\varepsilon_n, a_n}(t_i) = g^{\delta_n}_{\varepsilon_n}(t_i)$ on Ω_D. We have $w_n \to w := (u(t_1), \ldots, u(t_k))$ in $SBV(\Omega'; \mathbb{R}^k)$, where $u(t_i) = g(t_i)$ on Ω_D. Note that for all n we have $S_{w_n} = \bigcup_{i=1}^k S_{u^{\delta_n}_{\varepsilon_n, a_n}(t_i)}$ so that

$$\mathcal{H}^1(S_{w_n}) \leq \lambda_n(t).$$

Passing to the limit for $n \to +\infty$ and applying Ambrosio's Theorem we get

$$\mathcal{H}^1\left(\bigcup_{i=1}^k S_{u(t_i)}\right) = \mathcal{H}^1(S_w) \leq \liminf_n \mathcal{H}^1(S_{w_n}) \leq \lambda(t);$$

we thus have

$$\mathcal{H}^1\left(\bigcup_{i=1}^k S^{g(t_i)}(u(t_i))\right) = \mathcal{H}^1(S_w) \leq \lambda(t)$$

and taking the sup over all $t_1, \ldots, t_k$, we obtain (4.27) in D. The case $t \notin D$ follows since $\mathcal{H}^1(\Gamma(\cdot))$ is left continuous by definition.

Lemma 4.6. *For every $t \in [0,1]$ there exists $u(t) \in SBV(\Omega)$ such that the following hold:*

(a) *for all $t \in [0,1]$*

$$S^{g(t)}(u(t)) \subseteq \Gamma(t) \quad \text{up to a set of } \mathcal{H}^1\text{-measure } 0, \tag{4.28}$$

and for all $t \in [0,1]$ and for all $v \in SBV(\Omega)$

$$\|\nabla u(t)\|^2 \leq \|\nabla v\|^2 + \mathcal{H}^1\left(S^{g(t)}(v) \setminus \Gamma(t)\right); \tag{4.29}$$

(b) *∇u is continuous in $[0,1] \setminus (D \cup \mathcal{N})$ with respect to the strong topology of $L^2(\Omega; \mathbb{R}^2)$;*

(c) *if $\tilde{\mathcal{N}}$ is the set of discontinuities of the function η defined in* (4.26)*, for all $t \in [0,1] \setminus \tilde{\mathcal{N}}$ we have that*

$$\nabla u_{a_n}(t) \to \nabla u(t) \quad \text{strongly in } L^2(\Omega, \mathbb{R}^2).$$

Finally

$$\mathcal{E}(t) \geq \mathcal{E}(0) + 2\int_0^t \int_\Omega \nabla u(\tau) \nabla \dot{g}(\tau)\, dx\, d\tau, \tag{4.30}$$

where

$$\mathcal{E}(t) := \|\nabla u(t)\|^2 + \mathcal{H}^1(\Gamma(t)).$$

Proof. The definition of $u(t)$ is carried out as in Lemma 4.3 considering $t \in [0,1] \setminus D$, $t_n \in D$ with $t_n \nearrow t$, and the limit (up to a subsequence) of $u(t_n)$ in $SBV(\Omega)$: (4.28) and (4.29) hold, so that point (a) is proved. It turns out that $\nabla u(t)$ is uniquely determined and that it is left continuous in $[0,1] \setminus D$. Let us consider $t \in [0,1] \setminus (D \cup \mathcal{N})$, and let $t_n \searrow t$. By Ambrosio's Theorem, we have that there exists $u \in SBV(\Omega)$ with such that, up to a subsequence, $u(t_n) \to u$ in $SBV(\Omega)$. Since t is a continuity point of $\mathcal{H}^1(\Gamma(\cdot))$, we deduce that

$S^{g(t)}(u) \subseteq \Gamma(t)$ up to a set of $\mathcal{H}^1$-measure 0. Moreover by the minimality property for $u(t_n)$ and the fact $\Gamma(t) \subseteq \Gamma(t_n)$, we have that for all $v \in SBV(\Omega)$ with

$$\|\nabla u(t_n)\|^2 \leq \|\nabla v - \nabla g(t) + \nabla g(t_n)\|^2 + \mathcal{H}^1\left(S^{g(t)}(v) \setminus \Gamma(t_n)\right) \leq$$
$$\leq \|\nabla v - \nabla g(t) + \nabla g(t_n)\|^2 + \mathcal{H}^1\left(S^{g(t)}(v) \setminus \Gamma(t)\right),$$

and so we deduce that (4.29) holds with u in place of $u(t)$, and that $\nabla u(t_n) \to \nabla u$ strongly in $L^2(\Omega;\mathbb{R}^2)$. We obtain by uniqueness that $\nabla u = \nabla u(t)$, and so $\nabla u(\cdot)$ is continuous in $[0,1] \setminus (D \cup \mathcal{N})$ and this proves point (b). Point (c) follows in the same way of point (d) of Lemma 4.3.

Let us come to the proof of (4.30). Given $t \in [0,1]$ and $k > 0$, let $s_i^k := \frac{i}{k}t$ for all $i = 0, \ldots, k$. Let us set $u^k(s) := u(s_{i+1}^k)$ for $s_i^k < s \leq s_{i+1}^k$. By (4.29), comparing $u(s_i^k)$ with $u(s_{i+1}^k) - g(s_{i+1}^k) + g(s_i^k)$, it is easy to see that

$$\mathcal{E}(t) \geq \mathcal{E}(0) + 2\int_0^t \int_\Omega \nabla u^k(\tau)\nabla\dot{g}(\tau)\, d\tau\, dx + o_k,$$

where $o_k \to 0$ as $k \to +\infty$. Since ∇u is continuous with respect to the strong topology of $L^2(\Omega;\mathbb{R}^2)$ in $[0,1]$ up to a countable set, passing to the limit for $k \to +\infty$ we deduce (4.30). $\square$

We are now ready to prove the main result of the paper.

PROOF OF THEOREM 1.1. Let D be a countable and dense set in $[0,1]$ such that $0 \in D$, and let $(\delta_n, \varepsilon_n, a_n)$ and $\{t \to u(t) \in SBV(\Omega) : t \in [0,1]\}$ be the sequence and the evolution determined in Lemma 4.5 and Lemma 4.6. Let us set

$$u_n := u^{\delta_n}_{\varepsilon_n,a_n}, \qquad \Gamma_n := \Gamma^{\delta_n}_{\varepsilon_n,a_n}, \qquad \mathcal{E}_n := \mathcal{E}^{\delta_n}_{\varepsilon_n,a_n}.$$

Let $\overline{\mathcal{N}}$ be the union of the sets of discontinuities of η and $\mathcal{H}^1(\Gamma(\cdot))$, where η is defined in (4.26). Let $\mathcal{B} := \bigcap_{k=1}^{+\infty}\bigcup_{h=k}^{\infty}\mathcal{B}_h$, where $\mathcal{B}_h$ are as in Lemma 4.5; since $|\bigcup_{h=k}^{\infty}\mathcal{B}_h| < 2^{-k+1}$, we have $|\mathcal{B}| = 0$. For all $t \in [0,1] \setminus (\mathcal{B} \cup \overline{\mathcal{N}})$ we claim that

$$\nabla u_n(t) \to \nabla u(t) \quad \text{strongly in } L^2(\Omega;\mathbb{R}^2). \tag{4.31}$$

In fact, since $t \notin \bigcup_{h=k}^{\infty}\mathcal{B}_h$ for some k, by Lemma 4.5 we have

$$\lim_n \|\nabla u^{\delta_n}_{\varepsilon_n,a_n}(t) - \nabla u_{a_n}(t)\| = 0;$$

for $t \notin \overline{\mathcal{N}}$, by Lemma 4.6 we have that $\nabla u_{a_n}(t) \to \nabla u(t)$ strongly in $L^2(\Omega;\mathbb{R}^2)$ and so (4.31) holds.

Since $g_{\varepsilon_n} \to g$ strongly in $W^{1,1}([0,1];H^1(\Omega))$, we deduce that for a.e. $\tau \in [0,1]$

$$\nabla\dot{g}_{\varepsilon_n}(\tau) \to \nabla\dot{g}(\tau) \quad \text{strongly in } L^2(\Omega;\mathbb{R}^2).$$

Since $\mathcal{E}_n(0) \to \mathcal{E}(0)$ by (4.25) and $o^{\delta_n}_{\varepsilon_n} \to 0$, by semicontinuity of the energy and by (4.9) we have that for all $t \in D$

$$\mathcal{E}(t) \leq \liminf_n \mathcal{E}_n(t) \leq \limsup_n \mathcal{E}_n(t) \leq \mathcal{E}(0) + 2\int_0^t \int_\Omega \nabla u(\tau)\nabla\dot{g}(\tau)\, dx\, d\tau. \tag{4.32}$$

In view of (4.30), we conclude that for all $t \in D$

$$\mathcal{E}(t) = \mathcal{E}(0) + 2\int_0^t \int_\Omega \nabla u(\tau)\nabla\dot{g}(\tau)\, dx\, d\tau,$$

and since $\nabla u(\cdot)$ and $\mathcal{H}^1(\Gamma(\cdot))$ are left continuous at $t \notin D$ and so $\mathcal{E}(\cdot)$ is, we conclude that the equality holds for all $t \in [0,1]$. As a consequence $\{t \to u(t)\,, t \in [0,1]\}$ is a quasi-static evolution of brittle fractures. Let us prove that (4.32) is indeed true for all $t \in [0,1]$. In fact, if $t \notin D$, it is sufficient to prove

$$\liminf_n \mathcal{E}_n(t) \geq \mathcal{E}(t). \tag{4.33}$$

Considering $s \geq t$ with $s \in D$, by (4.7) we have

$$\mathcal{E}_n(s) \leq \mathcal{E}_n(t) + \int_{t^{\delta_n}_{j_n}}^{s^{\delta_n}_{j_n}} \int_\Omega \nabla u_n(\tau) \nabla \dot{g}_{\varepsilon_n}(\tau)\, dx\, d\tau + o^{\delta_n}_{\varepsilon_n} \quad t^{\delta_n}_{j_n} \leq t < t^{\delta_n}_{j_n+1},\ s^{\delta_n}_{j_n} \leq s < s^{\delta_n}_{j_n+1},$$

so that

$$\liminf_n \mathcal{E}_n(t) \geq \mathcal{E}(s) - \int_t^s \int_\Omega \nabla u(\tau) \nabla \dot{g}(\tau)\, dx\, d\tau.$$

Letting $s \searrow t$, since $\mathcal{E}(\cdot)$ is continuous, we have (4.33) holds. By (4.32) we deduce that $\mathcal{E}_n(t) \to \mathcal{E}(t)$ for all $t \in [0,1]$, so that point (b) is proved.

We now come to point (a). Since $\lambda(t) \geq \mathcal{H}^1(\Gamma(t))$ for all $t \in [0,1]$, by (4.31) and point (b), we deduce that $\lambda = \mathcal{H}^1(\Gamma(\cdot))$ in $[0,1]$ up to a set of measure 0. Since they are increasing functions, we conclude that λ and $\mathcal{H}^1(\Gamma(\cdot))$ share the same set of continuity points $[0,1] \setminus \mathcal{N}$, and that $\lambda = \mathcal{H}^1(\Gamma(\cdot))$ on $[0,1] \setminus \mathcal{N}$. In view of (4.31), point (a) is thus established for all t except $t \in (\mathcal{B} \cup \overline{\mathcal{N}}) \setminus \mathcal{N}$. In order to treat this case, we use the following argument. Considering the measures $\mu_n := \mathcal{H}^1 \llcorner \Gamma_n(t)$, we have that, up to a subsequence, $\mu_n \overset{*}{\rightharpoonup} \mu$ weakly-star in the sense of measures, and as a consequence of Ambrosio's Theorem we have $\mathcal{H}^1 \llcorner \Gamma(t) \leq \mu$ as measures. Since $t \notin \mathcal{N}$ we have $\mu_n(\mathbb{R}^2) \to \mathcal{H}^1(\Gamma(t))$, and so we deduce $\mathcal{H}^1 \llcorner \Gamma(t) = \mu$. Let us consider now $u_n(t)$; we have up to a subsequence $u_n(t) \to u$ in $SBV(\Omega)$ for some $u \in SBV(\Omega)$. Setting $u_n(t) := g^{\delta_n}_{\varepsilon_n}(t)$ and $u := g(t)$ on Ω_D, we have $u_n(t) \to u$ in $SBV(\Omega')$, and as a consequence of Ambrosio's Theorem, we get that $\mathcal{H}^1 \llcorner S^{g(t)}(u) \leq \mu = \mathcal{H}^1 \llcorner \Gamma(t)$, that is $S^{g(t)}(u) \subseteq \Gamma(t)$. By Theorem 2.3, we deduce that u is a minimum for

$$\min\{\|\nabla v\|^2 \,:\, S^{g(t)}(v) \subseteq \Gamma(t) \text{ up to a set of } \mathcal{H}^1\text{-measure } 0\ \},$$

and by uniqueness of the gradient we get that $\nabla u = \nabla u(t)$, so that the proof is concluded. ☐

5. Piecewise Affine Transfer of Jump and Proof of Proposition 4.2

The proof of Proposition 4.2 is based on the following proposition, which is a variant of Theorem 2.3 in the context of piecewise affine approximation.

Proposition 5.1. *Given $\varepsilon_n \to 0$, let $g^r_n \in H^1(\Omega)$ be such that $g^r_n \in \mathcal{AF}_{\varepsilon_n}(\Omega)$ and $g^r_n \to g^r$ strongly in $H^1(\Omega)$ for all $r = 0, \dots, i$. If $u^r_n \in \mathcal{A}_{\varepsilon_n,a}(\Omega)$ is such that $u^r_n \to u^r$ in $SBV(\Omega)$ for $r = 0, \dots, i$, then for all $v \in SBV(\Omega)$ with $\mathcal{H}^1\left(S^{g^i}(v)\right) < +\infty$ and $\nabla v \in L^2(\Omega;\mathbb{R}^2)$, there exists $v_n \in \mathcal{A}_{\varepsilon_n,a}(\Omega)$ such that $v_n \to v$ strongly in $L^1(\Omega)$, $\nabla v_n \to \nabla v$ strongly in $L^2(\Omega;\mathbb{R}^2)$ and*

$$\limsup_n \mathcal{H}^1\left(S^{g^i_n}(v_n) \setminus \bigcup_{r=0}^{i} S^{g^r_n}(u^r_n)\right) \leq \mu(a)\mathcal{H}^1\left(S^{g(t)}(v) \setminus \bigcup_{r=0}^{i} S^{g^r}(u^r)\right), \tag{5.1}$$

where $\mu :]0;\frac{1}{2}[\to \mathbb{R}$ with $\lim_{a\to 0^+} \mu(a) = 1$.

In view of Proposition 5.1, we can now prove Proposition 4.2.

Proof of Proposition 4.2. Notice that, in order to prove (4.14), it is sufficient to prove the existence of $\mu :]0; \frac{1}{2}[\to \mathbb{R}$ with $\lim_{a \to 0^+} \mu(a) = 1$ such that, given $t \in D$, for every $0 = t_0 \leq \ldots \leq t_r \leq \ldots \leq t_i = t$, $t_r \in D$, for all $v \in SBV(\Omega)$ we have

$$\|\nabla u_a(t)\|^2 \leq \|\nabla v\|^2 + \mu(a)\mathcal{H}^1 \left(S^{g(t)}(v) \setminus \bigcup_{r=0}^{i} S^{g(t_r)}(u_a(t_r)) \right). \tag{5.2}$$

In fact, taking the sup over all possible $t_0, \ldots, t_i$, we get (4.14).

We apply Proposition 5.1 considering $g_n^r := g_{\varepsilon_n}^{\delta_n}(t_r)$, $g^r := g(t_r)$, $u_n^r := u_{\varepsilon_n,a}^{\delta_n}(t_r)$, and $u^r := u_a(t_r)$ for $r = 0, \ldots, i$. There exists $\mu :]0; \frac{1}{2}[\to \mathbb{R}$ with $\lim_{a \to 0^+} \mu(a) = 1$ such that for $v \in SBV(\Omega)$, there exists $v_n \in \mathcal{A}_{\varepsilon_n,a}(\Omega)$ with $\nabla v_n \to \nabla v$ strongly in $L^2(\Omega; \mathbb{R}^2)$ and

$$\limsup_n \mathcal{H}^1 \left(S^{g_{\varepsilon_n}^{\delta_n}(t)}(v_n) \setminus \bigcup_{r=0}^{i} S^{g_{\varepsilon_n}^{\delta_n}(t_r)}(u_{\varepsilon_n,a}^{\delta_n}(t_r)) \right) \leq$$
$$\leq \mu(a)\mathcal{H}^1 \left(S^{g(t)}(v) \setminus \bigcup_{r=0}^{i} S^{g(t_r)}(u_a(t_r)) \right),$$

Comparing $u_{\varepsilon_n,a}^{\delta_n}(t)$ and v_n by means of (4.6), we obtain

$$\begin{aligned} \|\nabla u_{\varepsilon_n,a}^{\delta_n}(t)\|^2 &\leq \|\nabla v_n\|^2 + \mathcal{H}^1 \left(S^{g_{\varepsilon_n}^{\delta_n}(t)}(v_n) \setminus \Gamma_{\varepsilon_n,a}^{\delta_n}(t) \right) \leq \\ &\leq \|\nabla v_n\|^2 + \mathcal{H}^1 \left(S^{g_{\varepsilon_n}^{\delta_n}(t)}(v_n) \setminus \bigcup_{r=0}^{i} S^{g_{\varepsilon_n}^{\delta_n}(t_r)}(u_{\varepsilon_n,a}^{\delta_n}(t_r)) \right), \end{aligned} \tag{5.3}$$

so that, passing to the limit for $n \to +\infty$, we obtain that (5.2) holds. Moreover, we have that choosing $v = u_a(t)$, and taking the limsup in (5.3), we get that $\nabla u_{\varepsilon_n,a}^{\delta_n}(t) \to \nabla u_a(t)$ strongly in $L^2(\Omega; \mathbb{R}^2)$. ☐

The rest of the section is devoted to the proof of Proposition 5.1. It will be convenient, as in Section 4, to consider Ω_D polygonal open bounded subset of $\mathbb{R}^2$ such that $\Omega_D \cap \Omega = \emptyset$ and $\partial\Omega \cap \partial\Omega_D = \partial_D\Omega$ up to a finite number of vertices; we set $\Omega' := \Omega \cup \Omega_D \cup \partial_D\Omega$. We suppose that $\mathbf{R}_\varepsilon$ can be extended to a regular triangulation of Ω' which we still indicate by $\mathbf{R}_\varepsilon$.

We need several preliminary results. Let us set $z_n^r := u_n^r - g_n^r$, and let us extend z_n^r to zero on Ω_D. Similarly, we set $z^r := u^r - g^r$, and we extend z^r to zero on Ω_D.

Let $\sigma > 0$, and let C be the set of corners of $\partial_D\Omega$. Let us fix $G \subseteq \mathbb{R}$ countable and dense: we recall that for all $r = 0, \ldots, i$ we have up to a set of $\mathcal{H}^1$-measure zero

$$S_{z^r} = \bigcup_{c_1, c_2 \in G} \partial^* E_{c_1}(r) \cap \partial^* E_{c_2}(r),$$

where $E_c(r) := \{x \in \Omega' : z^r(x) > c\}$ and ∂^* denotes the essential boundary (see [4]). Let us consider

$$J_j := \{x \in \bigcup_{r=0}^{i} S_{z^r} \setminus C : (z^l)^+(x) - (z^l)^-(x) > \frac{1}{j} \text{ for some } l = 0, \ldots, i\},$$

with j so large that $\mathcal{H}^1(\bigcup_{r=0}^{i} S_{z^r} \setminus J_j) \leq \sigma$. Let U be a neighborhood of $\bigcup_{r=0}^{i} S_{z^r}$ such that $|U| \leq \frac{\sigma}{j^2}$. Following [10, Theorem 2.1] (see Fig.3), we can find a finite disjoint collection of closed cubes $\{Q_k\}_{k=1,\ldots,K}$ with center $x_k \in J_j$, edge of length $2r_k$ and oriented as the normal

$\nu(x_k)$ to $S_{z^{r(k)}}$ at x_k, such that $\bigcup_{k=1}^K Q_k \subseteq U$ and $\mathcal{H}^1(J_j \setminus \bigcup_{k=1}^K Q_k) \leq \sigma$. Moreover for all $k = 1, \ldots, K$ there exists $r(k) \in \{0, \ldots, i\}$ and $c_1(r(k)), c_2(r(k)) > 0$ such that

$$\mathcal{H}^1\left(\left[\bigcup_{r=0}^{i} S_{z^r} \setminus S_{z^{r(k)}}\right] \cap Q_k\right) \leq \sigma r_k,$$

and the following hold



Fig. 3

(a) if $x_k \in \Omega$ then $Q_k \subseteq \Omega$, and if $x_k \in \partial_D\Omega$ then $Q_k \cap \partial_D\Omega = H_k$, where H_k denotes the intersection of Q_k with the straight line through x_k orthogonal to $\nu(x_k)$;

(b) $\mathcal{H}^1(S_{z^{r(k)}} \cap \partial Q_k) = 0$;

(c) $r_k \leq c\mathcal{H}^1(S_{z^{r(k)}} \cap Q_k)$ for some $c > 0$;

(d) $(z^{r(k)})^-(x) < c_1(r(k)) < c_2(r(k)) < (z^{r(k)})^+(x)$ and $c_2(r(k)) - c_1(r(k)) \geq \frac{1}{2j}$;

(e) $\mathcal{H}^1([S_{z^{r(k)}} \setminus \partial^* E_{c_s(r(k))}(r(k))] \cap Q_k) \leq \sigma r_k$ for $s = 1, 2$;

(f) if $s = 1, 2$, $\mathcal{H}^1(\{y \in \partial^* E_{c_s(r(k))}(r(k)) \cap Q_k \,:\, \mathrm{dist}(y, H_k) \geq \frac{\sigma}{2} r_k\}) < \sigma r_k$;

(g) if $Q_k^+ := \{x \in Q_k \,|\, x \cdot \nu(x_k) > 0\}$ and $s = 1, 2$

$$\|1_{E_{c_s(r(k))}(r(k)) \cap Q_k} - 1_{Q_k^+}\|_{L^1(\Omega')} \leq \sigma^2 r_k^2; \tag{5.4}$$

(h) $\mathcal{H}^1((S_v \setminus S_{z^{r(k)}}) \cap Q_k) < \sigma r_k$ and $\mathcal{H}^1(S_v \cap \partial Q_k) = 0$.

Let us indicate by R_k the intersection of Q_k with the strip centered in H_k with width $2\sigma r_k$, and let us set $V_k^\pm := \{x_k \pm r_k e(x_k) + s\nu(x_k) \,:\, s \in \mathbb{R}\} \cap R_k$, where $e(x_k)$ is such that $\{e(x_k), \nu(x_k)\}$ is an orthonormal base of $\mathbb{R}^2$ with the same orientation of the canonical one.

For all $B \subseteq \Omega'$, let us set

$$\mathcal{R}_n(B) := \{T \in \mathbf{R}_{\varepsilon_n} \,:\, T \cap B \neq \emptyset\}, \qquad \mathcal{T}_n^k(B) := \{T \in \mathbf{T}(z_n^{r(k)}) \,:\, T \cap B \neq \emptyset\}.$$

In the following, we will often indicate with the same symbol a family of triangles and their support in $\mathbb{R}^2$, being clear from the context in which sense has to be intended. We will consider $z_n^{r(k)}$ defined pointwise in $\Omega' \setminus \overline{S}_{z_n^{r(k)}}$ and so the upper levels of $z_n^{r(k)}$ are intended as subsets of $\Omega' \setminus \overline{S}_{z_n^{r(k)}}$.

Lemma 5.2. *For all $k = 1, \dots, K$ there exists $c_n^k \in [c_1(r(k)), c_2(r(k))]$ such that, setting $E_n^k := \{x \in \mathcal{R}_n(Q_k) \,:\, z_n^{r(k)}(x) > c_n^k\}$, we have*

$$\limsup_n \sum_{k=1}^K \mathcal{H}^1\left((\partial_{\mathcal{R}_n(Q_k)} E_n^k) \setminus S_{z_n^{r(k)}} \right) = o_\sigma, \tag{5.5}$$

and

$$\limsup_n \| 1_{E_n^k} - 1_{Q_k^+} \|_{L^1(\Omega')} \leq \sigma^2 r_k^2, \tag{5.6}$$

where $\partial_{\mathcal{R}_n(Q_k)}$ denotes the boundary operator in $\mathcal{R}_n(Q_k)$, and $o_\sigma \to 0$ as $\sigma \to 0$.

Proof. Note that for n large we have $\bigcup_{k=1}^K \mathcal{R}_n(Q_k) \subseteq U$, so that $|\bigcup_{k=1}^K \mathcal{R}_n(Q_k)| \leq \frac{\sigma}{j^2}$. By Hölder inequality and since $\|\nabla z_n^r\| \leq C'$ for all $r = 0, \dots, i$, it follows that

$$\sum_{r=0}^i \int_{\{\cup_k \mathcal{R}_n(Q_k) : r(k) = r\}} |\nabla z_n^r| \, dx \leq \sum_{r=0}^i \|\nabla z_n^r\| \frac{\sqrt{\sigma}}{j} \leq (i+1) C' \frac{\sqrt{\sigma}}{j}.$$

Following [10, Theorem 2.1], we can apply coarea-formula for BV-functions (see [4]) taking into account that $z_n^{r(k)}$ belongs to $SBV(\Omega')$ so that the singular part of the derivative is carried only by $S_{z_n^{r(k)}}$: since for n large the $\mathcal{R}_n(Q_k)$'s are disjoint, we obtain

$$\sum_{k=1}^K \int_{\mathbb{R}} \mathcal{H}^1 \Big(\big(\partial E_{c,n}(r(k)) \cap \mathcal{R}_n(Q_k)\big) \setminus S_{z_n^{r(k)}} \Big) \, dc \leq (i+1) C' \frac{\sqrt{\sigma}}{j}, \tag{5.7}$$

where $E_{c,n}(r(k)) := \{x \in \Omega' \setminus \overline{S}_{z_n^{r(k)}} \,:\, z_n^{r(k)}(x) > c\}$, and so

$$\sum_{k=1}^K \int_{c_1(r(k))}^{c_2(r(k))} \mathcal{H}^1 \Big(\big(\partial E_{c,n}(r(k)) \cap \mathcal{R}_n(Q_k)\big) \setminus S_{z_n^{r(k)}} \Big) \, dc \leq (i+1) C' \frac{\sqrt{\sigma}}{j}.$$

Notice that we can use the topological boundary instead of the reduced boundary of $E_{c,n}(r(k))$ in (5.7) since $z_n^{r(k)}$ is piecewise affine, and so $\partial E_{c,n}(r(k)) \setminus \partial^* E_{c,n}(r(k)) \neq \emptyset$ just for a finite number of c's. By the Mean Value Theorem we have that there exist $c_n^k \in [c_1(r(k)), c_2(r(k))]$ such that

$$\sum_{k=1}^K \mathcal{H}^1 \Big(\big(\partial E_{c_n^k,n}(r(k)) \cap \mathcal{R}_n(Q_k)\big) \setminus S_{z_n^{r(k)}} \Big) \leq 2iC'\sqrt{\sigma},$$

and taking the limsup for $n \to +\infty$, we get (5.5). Let us come to (5.6). Since

$$E_{c_2(r(k)),n}(r(k)) \subseteq E_{c_n^k,n}(r(k)) \subseteq E_{c_1(r(k)),n}(r(k)),$$

by (5.4) we have that for n large

$$\|1_{E_{c^k_n,n}(r(k))\cap Q_k} - 1_{Q_k^+}\|_{L^1(\Omega')} \leq \sigma^2 r_k^2,$$

and so, since $|\mathcal{R}_n(Q_k)\setminus Q_k| \to 0$, we conclude that (5.6) holds. $\square$

Fix $k \in \{1,\dots,K\}$, and let us consider the family $\mathcal{T}_n^k(E_n^k)$. Let us modify this family in the following way. Let $T \in \mathcal{T}_n^k(E_n^k)$; we keep it if $|T\cap E_n^k| > \frac{1}{2}|T|$, and we erase it otherwise. Let $E_n^{k,+}$ be this new family of triangles, and let $E_n^{k,-}$ be its complement in $\mathcal{T}_n^k(\mathcal{R}_n(Q_k))$.

Lemma 5.3. *For all $k = 1,\dots,K$ we have*

$$\limsup_n \sum_{k=1}^K \mathcal{H}^1\left(\partial_{\mathcal{R}_n(Q_k)} E_n^{k,+} \setminus S_{z_n^{r(k)}}\right) = o_\sigma, \tag{5.8}$$

and

$$\limsup_n \|1_{E_n^{k,+}} - 1_{Q_k^+}\|_1 \leq 4\sigma^2 r_k^2, \tag{5.9}$$

where $o_\sigma \to 0$ as $\sigma \to 0$.

Proof. Let $T \in \mathcal{T}_n^k(E_n^k)$. Since $z_n^{r(k)}$ is affine on T, it follows that $T\cap E_n^k$ is either a triangle with at least two edges contained in the edges of T or a trapezoid with three edges contained in the edges of T. Let $l(T)$ be the edge inside T where $z_n^{r(k)} = c_n^k$, where c_n^k is the value determining E_n^k (we consider $l(T) = \emptyset$ if $\mathrm{int}(T) \subseteq E_n^k$). In the case $T \in E_n^{k,+}$ as in the case $T \in E_n^{k,-}$, since the angles of the triangles of $\mathbf{T}(z_n^{r(k)})$ are uniformly bounded away from 0 and from π, arguing as in Lemma 2.4, we deduce that keeping or erasing T, we increase $\partial_{\mathcal{R}_n(Q_k)} E_n^k$ of a quantity which is less than $c\mathcal{H}^1(l(T))$ with c independent of ε_n. Then we have

$$\sum_{k=1}^K \mathcal{H}^1(\partial_{\mathcal{R}_n(Q_k)} E_n^{k,+} \setminus \partial_{\mathcal{R}_n(Q_k)} E_n^k) \leq \sum_{k=1}^K \sum_{T\in\mathcal{T}_n^k(E_n^k)} c\mathcal{H}^1(l(T)) \leq c\sum_{k=1}^K \mathcal{H}^1(\partial_{\mathcal{R}_n(Q_k)} E_n^k \setminus S_{z_n^{r(k)}}),$$

so that taking the limsup for $n \to +\infty$ and in view of (5.5) we deduce that (5.8) holds.

Let us come to (5.9). Note that $|\mathcal{T}_n^k(\partial Q_k^+)| \to 0$ as $n \to +\infty$. Then if $A_n^{k,+} := \{T \in \mathbf{T}(z_n^{r(k)}) : T \subseteq \mathrm{int}(Q_k^+)\}$, for n large we have

$$|Q_k^+ \setminus E_n^{k,+}| \leq |A_n^{k,+}\setminus E_n^{k,+}| + |\mathcal{T}_n^k(\partial Q_k^+)| \leq 2|Q_k^+ \setminus E_n^k| + |\mathcal{T}_n^k(\partial Q_k^+)|,$$

where the last inequality follows by construction of $E_n^{k,+}$. Taking the limsup for $n \to +\infty$, in view of (5.6) we get

$$\limsup_n |Q_k^+ \setminus E_n^{k,+}| \leq 2\sigma^2 r_k^2.$$

The inequality $\limsup_n |E_n^{k,+}\setminus Q_k^+| \leq 2\sigma^2 r_k^2$ follows analogously. $\square$

For all $k = 1,\dots,K$ and $s \in \mathbb{R}$, let us set

$$H_k(s) := \{x + s\nu(x_k),\ x \in H_k\}.$$

Lemma 5.4. *There exist $s_n^+ \in]\frac{\sigma}{4}r_k, \frac{\sigma}{2}r_k[$ and $s_n^- \in]-\frac{\sigma}{2}r_k, -\frac{\sigma}{4}r_k[$ such that, setting $H_n^{k,+} := H_k(s_n^+)$ and $H_n^{k,-} := H_k(s_n^-)$ we have for n large enough*

$$\mathcal{H}^1(H_n^{k,+}\setminus E_n^{k,+}) \leq 20\sigma r_k, \quad \mathcal{H}^1(H_n^{k,-}\cap E_n^{k,+}) \leq 20\sigma r_k.$$

Proof. By (5.9) we can write for n large

$$\int_{\frac{\sigma}{4}r_k}^{\frac{\sigma}{2}r_k} \mathcal{H}^1(H_k(s) \setminus E_n^{k,+})\, ds \le 5\sigma^2 r_k^2,$$

so that we get $s_n^+ \in]\frac{\sigma}{4}r_k, \frac{\sigma}{2}r_k[$ with

$$\mathcal{H}^1(H_k(s_n^+) \setminus E_n^{k,+}) \le 20\sigma r_k.$$

Similarly we can reason for s_n^-. ☐

Let $S_n^{k,+}$ be the straight line containing $H_n^{k,+}$: up to replacing $H_n^{k,+}$ by the connected component of $S_n^{k,+} \cap \mathcal{R}_n(Q_k)$ to which it belongs, we may suppose that $H_n^{k,+} \setminus E_n^{k,+}$ is a finite union of segments l_j^+ with extremes A_j and B_j belonging to the edges of the triangles of $\mathcal{T}_n^k(\mathcal{R}_n(Q_k))$ such that for n large

$$\mathcal{H}^1(H_n^{k,+} \setminus E_n^{k,+}) = \mathcal{H}^1\left(\bigcup_{j=1}^m l_j^+\right) \le 20\sigma r_k.$$

By Lemma 2.4, for all j there exists a curve L_j^+ inside the edges of the triangles of $\mathcal{T}_n^k(\mathcal{R}_n(Q_k))$ joining A_j and B_j and such that

$$\mathcal{H}^1(L_j^+) \le c\mathcal{H}^1(l_j^+), \tag{5.10}$$

with c independent of ε_n. Let us set

$$\gamma_n^{k,+} := L_1^+ \cup B_1A_2 \cup L_2^+ \cup \cdots \cup B_{m-1}A_m \cup L_m^+.$$

Similarly, let us construct $\gamma_n^{k,-}$ relative to $H_n^{k,-} \cap E_n^+$. Note that for n large enough $\gamma_n^{k,+} \cap H_k(\sigma) = \emptyset$, $\gamma_n^{k,-} \cap H_k(-\sigma) = \emptyset$, and $\gamma_n^{k,+} \cap \gamma_n^{k,-} = \emptyset$. Let us consider the connected component $\mathcal{C}_k^+$ of $\mathcal{R}_n(Q_k) \setminus \gamma_n^{k,+}$ containing $H_k(\sigma)$. Similarly, let us consider the connected component $\mathcal{C}_k^-$ of $\mathcal{R}_n(Q_k) \setminus \gamma_n^{k,-}$ containing $H_k(-\sigma)$. For n large enough, by (5.10)

$$\mathcal{H}^1\left(\partial_{\mathcal{R}_n(Q_k)}\mathcal{C}_k^+ \setminus \bigcup_{i=1}^{m-1} B_iA_{i+1}\right) \le c\sum_{j=1}^m \mathcal{H}^1(l_j^+) \le 20c\sigma r_k. \tag{5.11}$$

A similar estimate holds for $\partial_{\mathcal{R}_n(Q_k)}\mathcal{C}_k^-$.

Let $\tilde{E}_n^{k,+}$ be the family of triangles obtained adding to $E_n^{k,+}$ those $T \in E_n^{k,-}$ such that $T \subseteq \mathcal{C}_k^+$, and subtracting those $T \in E_n^{k,+}$ such that $T \subseteq \mathcal{C}_k^-$. Let $\tilde{E}_n^{k,-}$ be the complement of $\tilde{E}_n^{k,+}$ in $\mathcal{T}_n^k(\mathcal{R}_n(Q_k))$.

We claim that there exists $C > 0$ independent of n such that for all $k = 1, \dots, K$ and for n large

$$\mathcal{H}^1\left(\partial_{\mathcal{R}_n(Q_k)}\tilde{E}_n^{k,+} \setminus \partial_{\mathcal{R}_n(Q_k)}E_n^{k,+}\right) \le C\sigma r_k. \tag{5.12}$$

In fact, let ζ be an edge of $\partial_{\mathcal{R}_n(Q_k)}\tilde{E}_n^{k,+} \setminus \partial_{\mathcal{R}_n(Q_k)}E_n^{k,+}$, that is ζ belongs to a triangle T that has been changed in the operation above described. Let us assume for instance that $T \in E_n^{k,-}$ and $T \subseteq \mathcal{C}_k^+$. If T' is such that $T \cap T' = \zeta$, then $T' \in E_n^{k,-}$: in fact if by contradiction $T' \in E_n^{k,+}$, then $T' \in \tilde{E}_n^{k,+}$ and so we would have $\zeta \notin \partial_{\mathcal{R}_n(Q_k)}\tilde{E}_n^{k,+}$ which is absurd. Similarly we get $T' \not\subseteq \mathcal{C}_k^+$. This means that $\zeta \subseteq \partial_{\mathcal{R}_n(Q_k)}\mathcal{C}_k^+$, and since the horizontal edges of $\gamma_n^{k,+}$ intersect by construction only elements of $E_n^{k,+}$, we deduce that $\zeta \subseteq \partial_{\mathcal{R}_n(Q_k)}\mathcal{C}_k^+ \setminus (\cup_{i=1}^m A_iB_i)$, and by (5.11) we conclude that (5.12) holds.

We can summarize the previous results as follows.

Lemma 5.5. *For all $k = 1, \dots, K$ there exist two families $\tilde{E}_n^{k,+}$ and $\tilde{E}_n^{k,-}$ of triangles with $\mathcal{T}_n^k(\mathcal{R}_n(Q_k)) = \tilde{E}_n^{k,+} \cup \tilde{E}_n^{k,-}$, $Q_k^+ \setminus R_k \subseteq \tilde{E}_n^{k,+}$ and $Q_k^- \setminus R_k \subseteq \tilde{E}_n^{k,-}$, and such that*

$$\limsup_n \sum_{k=1}^K \mathcal{H}^1\left(\partial_{\mathcal{R}_n(Q_k)}\tilde{E}_n^{k,+} \setminus S_{z_n^{r(k)}}\right) = o_\sigma, \tag{5.13}$$

where $o_\sigma \to 0$ as $\sigma \to 0$. Moreover, in the case $x_k \in \partial_D\Omega$, we can modify $\tilde{E}_n^{k,+}$ or $\tilde{E}_n^{k,-}$ in such a way that $\tilde{E}_n^{k,+} \subseteq \Omega$ or $\tilde{E}_n^{k,-} \subseteq \Omega$.

Proof. We have that (5.13) follows from (5.8) and (5.12), and the fact that $\sum_{k=1}^K r_k \le c$, with c independent of σ. Let us consider the case $x_k \in \partial_D\Omega$ with $Q_k^+ \setminus R_k \subseteq \Omega$ (the other case being similar). From (5.13) we have that for n large $\sum_{k=1}^K \mathcal{H}^1\left(\partial_{\mathcal{R}_n(Q_k)}\tilde{E}_n^{k,+} \cap Q_k^-\right) \le o_\sigma$ because $z_n^{r(k)} = R_{\varepsilon_n} g_{h_n}(r(k))$ on Q_k^- and so there are no jumps in Q_k^-. We can thus redefine $\tilde{E}_n^{k,+}$ subtracting those triangles that are in Q_k^- obtaining again (5.13). ☐

We are now in position to prove Proposition 5.1.

Proof of Proposition 5.1. We work in the context of Ω'. For all $v \in SBV(\Omega')$ with $v = g^i$ on Ω_D, $\mathcal{H}^1(S_v) < +\infty$ and $\nabla v \in L^2(\Omega'; \mathbb{R}^2)$, we have to construct $v_n \in SBV(\Omega')$ such that $v_n = g_n^i$ on Ω_D, $(v_n)_{|\Omega} \in \mathcal{A}_{\varepsilon_n,a}(\Omega)$, $v_n \to v$ strongly in $L^1(\Omega')$, $\nabla v_n \to \nabla v$ strongly in $L^2(\Omega'; \mathbb{R}^2)$ and

$$\limsup_n \mathcal{H}^1\left(S_{v_n} \setminus \bigcup_{r=0}^i S_{u_n^r}\right) \le \mu(a)\mathcal{H}^1\left(S_v \setminus \bigcup_{r=0}^i S_{u^r}\right), \tag{5.14}$$

where we suppose that u_n^r and u^r are extended to Ω' setting $u_n^r := g_n^r$, and $u^r := g^r$ on Ω_D respectively.

We set $v = g^i + w$, where $w \in SBV(\Omega')$ with $w = 0$ on Ω_D. By density, it is sufficient to consider the case $w \in L^\infty(\Omega')$. Up to reducing U, we may assume that $\|\nabla g^i\|_{L^2(U;\mathbb{R}^2)} < \sigma$ and $\|\nabla w\|_{L^2(U;\mathbb{R}^2)} < \sigma$. Let R_k' be a rectangle centered in x_k, oriented as R_k, and such that $\overline{R_k'} \subset \mathrm{int} R_k$ and $\mathcal{H}^1(S_w \cap (R_k \setminus R_k')) < \sigma r_k$. We claim that there exists $w_\sigma \in SBV(\Omega')$ with $w_\sigma = w$ on $\bigcup_{k=1}^K R_k'$ and $w_\sigma = 0$ in Ω_D such that

(1) $\|w - w_\sigma\| + \|\nabla w - \nabla w_\sigma\| \le \sigma$;

(2) $\mathcal{H}^1(S_{w_\sigma} \cap (Q_k \setminus R_k')) \le o_\sigma r_k$, with $o_\sigma \to 0$ as $\sigma \to 0$;

(3) $\mathcal{H}^1(S_{w_\sigma} \setminus \bigcup_{k=1}^K R_k) \le \mathcal{H}^1(S_w \setminus \bigcup_{k=1}^K R_k) + \sigma$;

(4) $S_{w_\sigma} \setminus \bigcup_{k=1}^K R_k$ is union of disjoint segments with closure contained in $\Omega \setminus \bigcup_{k=1}^K R_k$;

(5) w_σ is of class $W^{2,\infty}$ on $\Omega \setminus \left(\bigcup_{k=1}^K R_k \cup \overline{S_{w_\sigma}}\right)$.

In fact, by Proposition 2.7, there exists $w_m \in SBV(\Omega')$ with $w_m = 0$ in $\Omega' \setminus \overline{\Omega}$ such that $w_m \to w$ strongly in $L^2(\Omega')$, $\nabla w_m \to \nabla w$ strongly in $L^2(\Omega';\mathbb{R}^2)$, S_{w_m} is polyhedral with $\overline{S_{w_m}} \subseteq \Omega$, w_m is of class $W^{2,\infty}$ on $\Omega \setminus \left(\bigcup_{k=1}^K R_k \cup \overline{S_{w_m}}\right)$, and $\lim_m \mathcal{H}^1(A \cap S_{w_m}) = \mathcal{H}^1(A \cap S_w)$ for all A open subset of Ω' with $\mathcal{H}^1(\partial A \cap S_w) = 0$. It is not restrictive to assume that $\mathcal{H}^1(S_w \cap \partial R_k) = 0$ and $\mathcal{H}^1(S_{w_m} \cap \partial R_k) = 0$ for all m. Let ψ_k be a smooth function such that $0 \leq \psi_k \leq 1$, $\psi_k = 1$ on R'_k and $\psi_k = 0$ outside R_k. Setting $\psi := \sum_{k=1}^K \psi_k$, let us consider $\tilde{w}_m := \psi w + (1-\psi) w_m$. Note that $\tilde{w}_m \to w$ strongly in $L^2(\Omega')$, $\nabla \tilde{w}_m \to \nabla w$ strongly in $L^2(\Omega';\mathbb{R}^2)$, $\tilde{w}_m = 0$ in Ω_D. Moreover, by capacity arguments, we may assume that $S_{\tilde{w}_m} \setminus \bigcup_{k=1}^K R_k$ is a finite union of disjoint segments with closure contained in $\Omega \setminus \bigcup_{k=1}^K R_k$. Finally, for $m \to +\infty$, we have

$$\mathcal{H}^1(S_{\tilde{w}_m} \setminus \bigcup_{k=1}^K R_k) \to \mathcal{H}^1(S_w \setminus \bigcup_{k=1}^K R_k)),$$

$$\mathcal{H}^1(S_{\tilde{w}_m} \cap \bigcup_{k=1}^K (Q_k \setminus R_k)) \to \mathcal{H}^1(S_w \cap \bigcup_{k=1}^K (Q_k \setminus R_k))$$

and $\limsup_m \mathcal{H}^1(S_{\tilde{w}_m} \cap (R_k \setminus R'_k)) \leq 2\mathcal{H}^1(S_w \cap (R_k \setminus R'_k)) \leq 2\sigma r_k$. Then we can take $w_\sigma := \tilde{w}_m$ for m large enough.

Let $S_{w_\sigma} \setminus \bigcup_{k=1}^K Q_k := \bigcup_{j=1}^m l_j$, where, by capacity arguments, we can always assume that l_j are disjoint segments with closure contained in $\Omega \setminus \bigcup_{k=1}^K Q_k$. We define a triangulation $\mathbf{T}_n \in \mathcal{T}_{\varepsilon_n,a}(\Omega')$ specifying its adaptive vertices as follows. Let us consider the families $\mathcal{R}_n(Q_k)$ and $\mathcal{R}_n(l_j)$ for $k = 1, \dots, K$ and $j = 1, \dots, m$. Note that for n large enough, $\mathcal{R}_n(Q_{k_1}) \cap \mathcal{R}_n(Q_{k_2}) = \emptyset$ for $k_1 \neq k_2$, $\mathcal{R}_n(l_{j_1}) \cap \mathcal{R}_n(l_{j_2}) = \emptyset$ for $j_1 \neq j_2$, and $\mathcal{R}_n(Q_k) \cap \mathcal{R}_n(l_j) = \emptyset$ for every k, j. We consider inside the triangles of $\mathcal{R}_n(Q_k)$ the adaptive vertices of $\mathbf{T}(z_n^{r(k)})$. Passing to $\mathcal{R}_n(l_j)$, by density arguments it is not restrictive to assume that l_j does not pass through the vertices of $\mathbf{R}_{\varepsilon_n}$ and that its extremes belong to the edges of $\mathbf{R}_{\varepsilon_n}$. Let $\zeta := [x,y]$ be an edge of $\mathcal{R}_n(l_j)$ such that $l_j \cap \zeta = \{P\}$. Proceeding as in [12], we take as adaptive vertex of ζ the projection of P on $\{tx + (1-t)y : t \in [a,(1-a)]\}$. Connecting these adaptive vertices, we obtain an *interpolating* polyhedral curve $\tilde{l}_j$ with

$$\mathcal{H}^1(\tilde{l}_j) \leq \mu(a)\mathcal{H}^1(l_j), \tag{5.15}$$

where μ is an increasing function such that $\lim_{a \to 0} \mu(a) = 1$. Finally, in the remaining edges, we can consider any admissible adaptive vertex, for example the middle point.

Let us define $w_n \in SBV(\Omega')$ in the following way. For all Q_k, let w_n be equal to w_σ on $\mathcal{R}_n(Q_k) \setminus R_k$, equal to the reflection of $w_{\sigma|Q_k^+ \setminus R_k}$ with respect to $H_k(\sigma)$ on $\tilde{E}_n^{k,+} \cap R_k$ and equal to the reflection of $w_{\sigma|Q_k^- \setminus R_k}$ with respect to $H_k(-\sigma)$ on $\tilde{E}_n^{k,-} \cap R_k$, where $\tilde{E}_n^{k,\pm}$ are defined as in Lemma 5.5. On the other elements of $\mathbf{T}_n$, let us set $w_n = w_\sigma$. Notice that $w_n = 0$ on Ω_D and that inside each $\mathcal{R}_n(Q_k)$, all the discontinuities of w_n are contained in $\partial_{\mathcal{R}_n(Q_k)} \tilde{E}_n^{k,+} \cup V_k \cup P_{w_\sigma}^k$, where $P_{w_\sigma}^k$ is the union of the polyhedral jumps of w_σ in $\mathcal{R}_n(Q_k)$ and of their reflected version with respect to $H_k(\pm\sigma)$. By Lemma 5.5 and since $\sum_{k=1}^K \mathcal{H}^1(V_k \cup P_{w_\sigma}^k) \leq o_\sigma$ with $o_\sigma \to 0$ as

$\sigma \to 0$, and $\mathcal{H}^1\left(\bigcup_{r=0}^{i} S_{z^r} \setminus \bigcup_{k=1}^{K} Q_k\right) \leq 2\sigma$, we have that

$$\begin{aligned}
&\limsup_n \mathcal{H}^1\left(S_{w_n} \setminus \bigcup_{r=0}^{i} S_{z_n^r}\right) \leq \\
&\leq \mathcal{H}^1\left(S_{w_\sigma} \setminus \bigcup_{k=1}^{K} Q_k\right) + \limsup_n \mathcal{H}^1\left((S_{w_n} \setminus \bigcup_{r=0}^{i} S_{z_n^r}) \cap \mathcal{R}_n(Q_k)\right) \leq \\
&\leq \mathcal{H}^1\left(S_{w_\sigma} \setminus \bigcup_{r=0}^{i} S_{z^r}\right) + \mathcal{H}^1\left(\bigcup_{r=0}^{i} S_{z^r} \setminus \bigcup_{k=1}^{K} Q_k\right) + o_\sigma \leq \\
&\leq \mathcal{H}^1\left(S_{w_\sigma} \setminus \bigcup_{r=0}^{i} S_{z^r}\right) + o_\sigma,
\end{aligned}$$

and since $\|\nabla w_\sigma\|_{L^2(U;\mathbb{R}^2)} \leq o_\sigma$ we get for n large

$$\|\nabla w_n\|^2_{L^2(\bigcup_{k=1}^{K} \mathcal{R}_n(Q_k))} \leq o_\sigma. \tag{5.16}$$

We now want to define an interpolation $\tilde{w}_n$ of w_n on $\mathbf{T}_n$. Firstly, we set $\tilde{w}_n = 0$ on all regular triangles of Ω_D. Passing to the triangles in $\mathcal{R}_n(Q_k)$ (see fig.4), by Lemma 2.5, we know that for n large enough, we have

$$\mathcal{H}^1(\partial\mathcal{R}_n(V_k)) \leq c\mathcal{H}^1(V_k), \quad \mathcal{H}^1(\partial\mathcal{R}_n(P^k_{w_\sigma})) \leq c\mathcal{H}^1(P^k_{w_\sigma}),$$

with c independent of n. If $T \in \mathcal{R}_n(V_k) \cup \mathcal{R}_n(P^k_{w_\sigma})$, we set $\tilde{w}_n = 0$ on T; otherwise, we define $\tilde{w}_n$ on T as the affine interpolation of w_n.

Since $\nabla\tilde{w}_n$ is uniformly bounded on $\mathcal{R}_n(H_k(\pm\sigma))$, $|\mathcal{R}_n(H_k(\pm\sigma))| \to 0$ and since w_n is uniformly bounded in $W^{2,\infty}$ on the triangles contained in $\mathcal{R}_n(Q_k) \setminus \mathcal{R}_n(V_k \cup P^k_{w_\sigma} \cup H_k(\pm\sigma))$ we have by the interpolation estimate (2.3) and by (5.16)

$$\limsup_n \|\nabla\tilde{w}_n\|^2_{L^2(\bigcup_{k=1}^{K} \mathcal{R}_n(Q_k))} \leq o_\sigma. \tag{5.17}$$

Moreover we have

$$\limsup_n \sum_{k=1}^{K} \mathcal{H}^1\left(\left(S_{\tilde{w}_n} \setminus \bigcup_{r=0}^{k} S_{z_n^r}\right) \cap \mathcal{R}_n(Q_k)\right) \leq o_\sigma. \tag{5.18}$$

Let us come to the triangles not belonging to $\mathcal{R}_n(Q_k)$ for $k = 1, \dots, K$. For all $j = 1, \dots, m$, we denote by $\hat{\mathcal{R}}_n(l_j)$ the family of regular triangles that have edges in common with triangles of $\mathcal{R}_n(l_j)$. For n large we have that $\hat{\mathcal{R}}_n(l_{j_1}) \cap \hat{\mathcal{R}}_n(l_{j_2}) = \emptyset$ for $j_1 \neq j_2$. On every regular triangle $T \notin \bigcup_{k=1}^{K} \mathcal{R}_n(Q_k) \cup \bigcup_{j=1}^{m} \hat{\mathcal{R}}_n(l_j)$, we define $\tilde{w}_n$ as the affine interpolation of w_σ. Since w_σ is of class $W^{2,\infty}$ on T and T is regular, we obtain by the interpolation estimate (2.3)

$$\|\tilde{w}_n - w_\sigma\|^2_{W^{1,2}(T)} \leq K\varepsilon_n \|w_\sigma\|_{W^{2,\infty}}. \tag{5.19}$$

Let us consider now those triangles that are contained in the elements of $\bigcup_{j=1}^{m} \hat{\mathcal{R}}_n(l_j)$. Following [12], we can define $\tilde{w}_n$ on every T in such a way that $\tilde{w}_n$ admits discontinuities only on $\tilde{l}_j$, and $\|\nabla\tilde{w}_n\|_{L^\infty(T)} \leq \|\nabla w_\sigma\|_\infty$. Since $|\hat{\mathcal{R}}_n(l_j)| \to 0$ as $n \to \infty$, we deduce that

$$\lim_n \|\nabla\tilde{w}_n\|^2_{L^2(\hat{\mathcal{R}}_n(l_j))} = 0. \tag{5.20}$$

Moreover by (5.15) and since $\mathcal{H}^1\left(\bigcup_{r=0}^{i} S_{z^r} \setminus \bigcup_{k=1}^{K} Q_k\right) \leq 2\sigma$, we have

$$\mathcal{H}^1\left(S_{\tilde{w}_n} \cap \bigcup_{j=1}^{m} \hat{\mathcal{R}}_n(l_j)\right) \leq \mu(a)\mathcal{H}^1\left(S_{w_\sigma} \setminus \bigcup_{i=1}^{k} S_{z^r}\right) + o_\sigma, \tag{5.21}$$

where $o_\sigma \to 0$ as $\sigma \to 0$.



Fig. 4

We are now ready to conclude. Let us consider $\hat{w}_n \in \mathcal{A}_{\varepsilon_n,a}(\Omega)$ defined as $\hat{w}_n := g_n^i + \tilde{w}_n$. We have $\hat{w}_n \to g^i + w_\sigma$ strongly in $L^2(\Omega')$. By (5.17), (5.19), (5.20) we get

$$\limsup_n \|\nabla \hat{w}_n\|^2 \leq \|\nabla g^i + \nabla w_\sigma\|^2 + o_\sigma,$$

while by (5.18) and (5.21) we have

$$\limsup_n \mathcal{H}^1\left(S_{\hat{w}_n} \setminus \bigcup_{r=0}^{i} S_{z_n^r}\right) \leq \mu(a)\mathcal{H}^1\left(S_{w_\sigma} \setminus \bigcup_{r=0}^{i} S_{z^r}\right) + o_\sigma.$$

Letting now $\sigma \to 0$, using a diagonal argument, we conclude that Proposition 5.1 holds. ☐

6. Revisiting the approximation by Francfort and Larsen

In this section we show how the arguments of Section 4 may be used to deal with the discrete in time approximation of quasi-static growth of brittle fractures proposed by Francfort and Larsen in [10]. More precisely, we prove that there is strong convergence of the gradient of the displacement (in particular convergence of the bulk energy) and convergence of the surface energy at all times of continuity of the length of the crack; moreover there is convergence of the total energy at any time.

We briefly recall the notation employed in [10]. Let I_∞ be countable and dense in $[0,1]$, and let $I_n := \{0 = t_0^n \leq \ldots \leq t_n^n = 1\}$ such that (I_n) is an increasing sequence of sets whose union is I_∞. Let $\Omega \subseteq \mathbb{R}^N$ be a Lipschitz bounded domain, and let $\partial\Omega = \partial\Omega_f^c \cup \partial\Omega_f$, where $\partial\Omega_f^c$ is open in the relative topology. Let $\Omega' \subseteq \mathbb{R}^N$ be open and such that $\overline{\Omega} \subseteq \Omega'$, and let $g \in W^{1,1}([0,1]; H^1(\Omega'))$. At any time t_k^n, Francfort and Larsen consider u_k^n minimizer of

$$\int_\Omega |\nabla v|^2\, dx + \mathcal{H}^{N-1}\left(S_v \setminus \left[\bigcup_{0\leq j\leq k-1} S_{u_j^n} \cup \partial\Omega_f\right]\right)$$

in $\{v \in SBV(\Omega') : v = g(t_k^n) \text{ in } \Omega' \setminus \overline{\Omega}\}$. Setting $u^n(t) := u_k^n$ for $t \in [t_k^n, t_{k+1}^n[$, and $\Gamma^n(t) := \bigcup_{s\leq t, s\in I_n} S_{u^n(s)} \cup \partial\Omega_f$, they prove that

$$\mathcal{E}^n(t) \leq \mathcal{E}^n(0) + 2\int_0^{t_k^n}\int_\Omega \nabla u^n(\tau)\nabla\dot{g}(\tau)\, dx\, d\tau + o_n, \qquad t \in [t_k^n, t_{k+1}^n[, \tag{6.1}$$

where $\mathcal{E}^n(t) := \int_\Omega |\nabla u^n(t)|^2\, dx + \mathcal{H}^{N-1}(\Gamma^n(t))$ and $o_n \to 0$ as $n \to +\infty$. Using Theorem 2.3, they obtain a subsequence of $(u^n(\cdot))$, still denoted by the same symbol, such that $u^n(t) \to u(t)$ in $SBV(\Omega')$ and $\nabla u^n(t) \to \nabla u(t)$ strongly in $L^2(\Omega'; \mathbb{R}^N)$ for all $t \in I_\infty$, with $u(t)$ a minimizer of

$$\int_\Omega |\nabla v|^2\, dx + \mathcal{H}^{N-1}(S_v \setminus \Gamma(t)),$$

where $\Gamma(t) := \bigcup_{s\in I_\infty, s\leq t} S_{u(s)} \cup \partial\Omega_f$. The evolution $\{t \to u(t),\ t \in I_\infty\}$ is extended to the whole $[0,1]$ using the approximation from the left in time.

We can now use the arguments of Section 4. Following Lemma 4.6, it turns out that for all $t \in [0,1]$

$$\mathcal{E}(t) \geq \mathcal{E}(0) + 2\int_0^t\int_\Omega \nabla u(\tau)\nabla\dot{g}(\tau)\, dx\, d\tau. \tag{6.2}$$

Moreover, by the Transfer of Jump and the uniqueness argument of Lemma 4.3, we have that $\nabla u^n(t) \to \nabla u(t)$ strongly in $L^2(\Omega'; \mathbb{R}^N)$ for all $t \notin \mathcal{N}$, where $\mathcal{N}$ is the (at most countable) set of discontinuities of the pointwise limit λ of $\mathcal{H}^{N-1}(\Gamma(\cdot))$ (which exists up to a further subsequence by Helly's Theorem). Then we pass to the limit in (6.1) obtaining

$$\mathcal{E}(t) \leq \mathcal{E}(0) + 2\int_0^t\int_\Omega \nabla u(\tau)\nabla\dot{g}(\tau)\, dx\, d\tau;$$

moreover, following the proof of Theorem 1.1, we have that for all $t \in [0,1]$

$$\mathcal{E}(t) \leq \liminf_n \mathcal{E}_n(t) \leq \limsup_n \mathcal{E}_n(t) = \mathcal{E}(0) + 2\int_0^t\int_\Omega \nabla u(\tau)\nabla\dot{g}(\tau)\, dx\, d\tau,$$

and taking into account (6.2) we get the convergence of the total energy at any time. Since $\nabla u^n(t) \to \nabla u(t)$ strongly in $L^2(\Omega'; \mathbb{R}^N)$ for every $t \in I_\infty$, we deduce that $\lambda = \mathcal{H}^{N-1}(\Gamma(\cdot))$ on I_∞, so that the convergence of the surface energy holds in I_∞. The extension to the continuity times for $\mathcal{H}^{N-1}(\Gamma(\cdot))$ follows like in the final part of the proof of Theorem 1.1.

ACKNOWLEDGMENTS

The authors wish to thank Gianni Dal Maso for having proposed them the problem, and Gianni Dal Maso and Gilles A. Francfort for many helpful and interesting discussions.

References

[1] Ambrosio L.: A compactness theorem for a new class of functions of bounded variations. *Boll. Un. Mat. Ital.* **3-B** (1989), 857-881.

[2] Ambrosio L.: Existence theory for a new class of variational problems. *Arch. Ration. Mech. Anal.* **111** (1990), 291-322.

[3] Ambrosio L.: A new proof of the SBV compactness theorem. *Calc. Var.* **3** (1995), 127-137.

[4] Ambrosio L., Fusco N., Pallara D.: *Functions of bounded variations and Free Discontinuity Problems.* Clarendon Press, Oxford, 2000.

[5] Bourdin B., Francfort G.A., Marigo J.J.: Numerical experiments in revisited brittle fracture. *J. Mech. Phys. Solids*, **48-4** (2000), 797-826.

[6] Brezis, H.: *Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert.* North-Holland, Amsterdam, 1973.

[7] Ciarlet P.G.: *The Finite Element Method for Elliptic Problems*, North Holland, Amsterdam (1978).

[8] Cortesani G.: Strong approximation of *GSBV* functions by piecewise smooth functions. *Ann. Univ. Ferrara - Sez VII - Sc. Mat.* **43** (1997), 27-49.

[9] Dal Maso G., Toader R.: A model for the quasi-static growth of brittle fractures: existence and approximation results. *Arch. Ration. Mech. Anal.* **162** (2002), 101-135.

[10] Francfort G.A., Larsen C.J.: Existence and convergence for quasi-static evolution in brittle fracture. *To appear.*

[11] Francfort G.A., Marigo J.-J.: Revisiting brittle fractures as an energy minimization problem. *J. Mech. Phys. Solids* **46** (1998), 1319-1342.

[12] Negri M.: Numerical methods for free-discontinuity problems based on approximations by Γ-convergence. PhD Thesis, SISSA/ISAS, (2001).

[13] Rogers C.A.:*Hausdorff Measures.* Cambridge University Press, Cambridge, 1970.

(Alessandro Giacomini) S.I.S.S.A., Via Beirut 2-4, 34014, Trieste, Italy
E-mail address, A. Giacomini: `giacomin@sissa.it`

(Marcello Ponsiglione) S.I.S.S.A., Via Beirut 2-4, 34014, Trieste, Italy
E-mail address, M. Ponsiglione: `ponsigli@sissa.it`